HEALTHCARE COST AND UTILIZATION PROJECT — HCUP A FEDERAL-STATE-INDUSTRY PARTNERSHIP IN HEALTH DATA Sponsored by the Agency for Healthcare Research and Quality

INTRODUCTION TO

THE HCUP NATIONWIDE READMISSIONS DATABASE (NRD)

2019

Please read all documentation carefully.

THE NRD CONTAINS A FULL YEAR OF INTERNATIONAL CLASSIFICATION OF DISEASES, TENTH REVISION, CLINICAL MODIFICATION/PROCEDURE CODING SYSTEM (ICD-10-CM/PCS) CODES BEGINNING DATA YEAR 2016.

These pages provide an introduction to the 2019 NRD.

For full documentation and notification of changes, visit the HCUP User Support (HCUP-US) website at www.hcup-us.ahrq.gov.

Issued October 2021

Agency for Healthcare Research and Quality Healthcare Cost and Utilization Project (HCUP) 5600 Fishers Lane Mail Stop 7W25B Rockville, MD 20857

NRD Data and Documentation Distributed through the HCUP Central Distributor:

Website: www.hcup-us.ahrq.gov Phone: (866) 290-4287 (toll-free) Fax: (805) 979-3787 Email: hcup@ahrq.gov

TABLE OF CONTENTS

| SUMMARY OF DATA USE Restrictions | 1 |
|--|----------------------------|
| HCUP DATA USE AGREEMENT REQUIREMENTS | 2 |
| HCUP CONTACT INFORMATION | 2 |
| HCUP Central Distributor and HCUP User Support | 2 |
| WHAT'S NEW IN THE 2019 NRD? | 4 |
| UNDERSTANDING THE NRD | 4 |
| ABSTRACT | 5 |
| INTRODUCTION TO THE NATIONWIDE READMISSIONS DATABASE (NRD) | 7 |
| Overview of NRD Data | 7 7 8 8 9 |
| ICD-10-CM/PCS Started October 1, 2015 at the Beginning of Fiscal Year 2016 | |
| File Structure of the NRD NRD Data Elements | |
| Getting Started Decompressing the NRD Files Downloading and Running the Load Programs NRD Documentation HCUP Online Tutorials HCUP Reports on Readmission Methodology | 14 14 15 15 15 |
| ICD-9-CM and ICD-10-CM/PCS Diagnosis and Procedure Codes | |
| SAMPLING DESIGN OF THE NRD | |
| Target Universe | |
| Sampling Frame | |
| Discharge Weights Post-Stratification for Weighting Weighting | 17 |
| Final NRD Sample | 18 |
| Limitations of the NRD | 19 |
| HOW TO USE THE NRD FOR READMISSION ANALYSES | 20 |
| NRD Data Elements Critical to Tracking a Patient and Determining the Time Between | 04 |
| Admissions Patient Linkage Number (NRD_VisitLink) Time Between Admissions (NRD_DaysToEvent and LOS) | 21 |
| Defining Readmissions | 22 |

| Defining the Index Event Specifying the Criteria for a Readmission Selecting the Time Period for Revisits Reporting Rates of Readmission | |
|---|----------------------------------|
| Calculating Nationally Weighted Estimates | 25 |
| Variance Calculations | 25 |
| Computer Software for Weighted and Variance Calcu | Ilations 26 |
| NRD READMISSION RATES REPORTED ON THE HCL | |
| APPENDIX A: NRD INTRODUCTORY INFORMATION | 1 |
| Table A.1. HCUP Partners Participating in the 2019 | VRD 2 |
| Figure A.1. HCUP States Participating in the 2019 N | २D 3 |
| Table A.2. Percentage of SID Discharges in the NRD | by Type of Discharge, 20191 |
| Table A.3. Summary of NRD States, Hospitals, and I | npatient Stays 1 |
| Table A.4. NRD-Related Reports and Database Doc | umentation Available on HCUP-US4 |
| Table A.5 Range of Discharge Weights by Patient Ag | je and Sex, 20195 |
| Table A.6 Rate and Number of 30-Day All-Cause Re Patient Age Group, 2019 | |
| Table A.7 Rate and Number of 30-Day All-Cause Re Principal Diagnoses at Index Admission, Ranked by | |
| Table A.8 Rate and Number of 30-Day All-Cause Re (MDC) at Index Admission, Ranked by Number of Re | |
| APPENDIX B: HANDLING OF TRANSFERS AND SAME | -DAY STAYS 1 |
| Defining Transfers and Same-Day Events | 2 |
| Creating a Combined Transfer Record | 3 |
| APPENDIX C: STATE-SPECIFIC RESTRICTIONS | 1 |
| Table C.1. State-Specific Restrictions | 2 |
| APPENDIX D: NRD DATA ELEMENTS AND CODES | 1 |
| Table D.1. Data Elements in the 2019 NRD Core File | 2 |
| Table D.2. Data Elements in the 2019 NRD Severity | Measures File6 |
| Table D.3. Data Elements in the 2019 NRD Diagnosi | s and Procedure Groups Files7 |
| Table D.4. Data Elements in the 2019 NRD Hospital | File 9 |
| APPENDIX E: EVALUATION OF THE DIFFERENCE IN HCUP PATIENT LINKAGE NUMBERS BEING SPECIFIC | |
| Table E.1. Twenty Conditions with the Largest Numb Readmissions to Within State and Across All States - | |
| Table E.2. Ten Conditions with the Largest Difference Readmissions to Within State and Across All States | |

| Table E.3. Readmission Rates Restricting Readmissions to Within State and Across All | |
|--|---|
| States, Medicare Standard Analytic File, 2011 | 6 |

HCUP NATIONWIDE READMISSIONS DATABASE (NRD) SUMMARY OF DATA USE RESTRICTIONS

***** REMINDER *****

All users of the NRD must take the online HCUP Data Use Agreement (DUA) training course, and read and sign a Data Use Agreement. Details and links may be found on the following page.

Authorized users of HCUP data agree to the following restrictions:^a

- Will not use the data for any purpose other than research, analysis, and aggregate statistical reporting.
- Will not re-release any data to unauthorized users.
- Will not redistribute HCUP data by posting on any website or publishing in any other publicly accessible online repository. If a journal or publication requests access to data or analytic files, will cite restrictions on data sharing in the Data Use Agreement and direct them to AHRQ HCUP (www.hcup-us.ahrq.gov) for more information on accessing HCUP data.
- Will not identify or attempt to identify any individual, including by the use of vulnerability analysis or penetration testing. Methods that could be used to identify individuals directly or indirectly shall not be disclosed or published.
- Will not report any statistics where the number of observations (i.e., individual discharge records) in any given cell of tabulated data is less than or equal to 10 (≤10).
- Will not publish information that could identify individual establishments (e.g., hospitals) and will not contact establishments.
- Will not use the data concerning individual establishments for commercial or competitive purposes affecting establishments, or to determine rights, benefits, or privileges of establishments.
- Will not use the data for criminal and civil litigation, including expert witness testimony or for law enforcement activities.
- Will not use data elements from the proprietary severity adjustment software packages (e.g., 3M[™] APR-DRGs) for any commercial purpose or to disassemble, decompile, or otherwise reverse engineer the proprietary software.
- Will acknowledge in reports that data from the "Healthcare Cost and Utilization Project (HCUP)" were used, including names of the specific databases used for analysis.^b

Any violation of the limitations in the Data Use Agreement is punishable under Federal law by a fine, up to five years in prison, or both. Violations may also be subject to penalties under State statutes.

^a This is a summary of key terms of the Data Use Agreement for Nationwide Databases; please refer to the DUA for full terms and conditions.

^b Suggested citations for the HCUP databases are provided in the Requirements for Publishing with HCUP Data available at <u>www.hcup-us.ahrq.gov/db/publishing.jsp</u>.

HCUP DATA USE AGREEMENT REQUIREMENTS

All HCUP data users, including data purchasers and collaborators, must complete the online HCUP Data Use Agreement (DUA) Training Tool, and read and sign the HCUP Data Use Agreement. Proof of training completion and signed Data Use Agreements must be submitted to the HCUP Central Distributor as described below.

Data purchasers will be required to provide their DUA training completion code and will execute their DUAs electronically as a part of the online ordering process. The DUAs and training certificates for collaborators and others with access to HCUP data should be submitted directly to the HCUP Central Distributor using the contact information below.

The on-line DUA training course is available at: www.hcup-us.ahrq.gov/tech_assist/dua.jsp.

The **HCUP Nationwide Data Use Agreement** is available on the AHRQ-sponsored HCUP User Support (HCUP-US) website at: <u>https://www.hcup-us.ahrq.gov/team/NationwideDUA.pdf</u>.

HCUP CONTACT INFORMATION

HCUP Central Distributor and HCUP User Support

Information about the content of the HCUP databases is available on the HCUP User Support (HCUP-US) website (<u>www.hcup-us.ahrq.gov</u>).

If you have questions, please review the HCUP Frequently Asked Questions located at <u>www.hcup-us.ahrq.gov/tech_assist/faq.jsp</u>.

If you need further technical assistance, please contact the HCUP Central Distributor and User Support team at:

Phone: 866-290-HCUP (4287) (toll free) Email: <u>hcup@ahrq.gov</u> Fax: 866-792-5313 (toll free in the United States)

Mailing address: HCUP Central Distributor IBM Watson Health 5425 Hollister Avenue, Suite 140 Santa Barbara, CA 93111

WHAT IS THE NATIONWIDE READMISSIONS DATABASE (NRD)?

- The Nationwide Readmissions Database (NRD) is a database of all-payer hospital inpatient stays that can be used to generate national estimates of readmissions. Outcomes of interest include national readmission rates, reasons for returning to the hospital for care, and the hospital costs for discharges with and without readmissions.
- The NRD addresses a large gap in healthcare data—the lack of nationally representative information on hospital readmissions for all patients, regardless of the expected payer for the hospital stay. The NRD is designed to be flexible to various types of analyses of readmissions. Criteria to determine the relationship between multiple hospital admissions for an individual patient in a calendar year are left to the analyst using the NRD.
- The NRD is drawn from HCUP State Inpatient Databases (SID) containing verified patient linkage numbers that can be used to track a person across hospitals within a State, while adhering to strict privacy guidelines. Unweighted, the NRD contains data from approximately 18 million discharges each year. Weighted, it estimates roughly 35 million discharges in the United States.
- There are 30 HCUP Partner States that contributed to the 2019 NRD: Alaska, Arkansas, California, Colorado, Delaware, Florida, Georgia, Hawaii, Indiana, Iowa, Louisiana, Maryland, Massachusetts, Mississippi, Missouri, Nebraska, New Mexico, Nevada, New York, Oregon, Pennsylvania, South Carolina, South Dakota, Tennessee, Utah, Virginia, Vermont, Washington, Wisconsin, and Wyoming. These States are geographically dispersed and account for 61.8 percent of the total U.S. resident population and 60.4 percent of all U.S. hospitalizations.
- The NRD is a publicly available database that can be purchased through the HCUP Central Distributor. The NRD is available for data years 2010–2019.
- Users must complete the <u>HCUP Data Use Agreement Training Course</u> prior to receiving the data.

WHAT'S NEW IN THE 2019 NRD?

- Data elements indicating certain types of hospital stays, including births, deliveries, injuries, hospital service lines, and operating room procedures are available in the NRD Core file.
- Data elements derived from the Elixhauser Comorbidity Software Refined for ICD-10-CM, the Clinical Classifications Software Refined (CCSR) for ICD-10-PCS procedures, and Procedure Classes Refined for ICD-10-PCS are available in the NRD Diagnosis and Procedure Groups file.

UNDERSTANDING THE NRD

- This document, *Introduction to the NRD, 2019,* summarizes the content of the 2019 NRD and describes the development of the NRD sample and weights. Similar documents for previous years of the NRD are available on the HCUP User Support (HCUP-US) website at www.hcup-us.ahrq.gov/db/nation/nrd/nrdarchive.jsp.
- In-depth documentation for the NRD is available at <u>www.hcup-us.ahrq.gov/db/nation/nrd/nrddbdocumentation.jsp</u>. Please refer to detailed documentation before using the data.
- Important considerations for data analysis are highlighted and references to further resources are provided.
- The coding system used to report diagnoses and procedures has changed over time:
 - Beginning with data year 2016, the NRD includes a full calendar year of data with diagnosis and procedure codes reported ICD-10-CM/PCS diagnosis and procedure codes.
 - In data year 2015, the first 9 months of the NRD contain International Classification of Diseases, Ninth Revision, Clinical Modification (ICD-9-CM) codes and the last 3 months contain ICD-10-CM/PCS codes.
 - In data year 2014 and prior years, the NRD contains ICD-9-CM diagnosis and procedure codes.
- The HCUP-US website has a section on <u>ICD-10-CM/PCS Resources</u> that summarizes key issues for researchers using HCUP and other administrative databases that include ICD-10-CM/PCS coding. The web page provides general guidance and forewarning to users analyzing outcomes that may be affected by the transition to the ICD-10-CM/PCS coding system and lists other related web resources.

HEALTHCARE COST AND UTILIZATION PROJECT — HCUP A FEDERAL-STATE-INDUSTRY PARTNERSHIP IN HEALTH DATA Sponsored by the Agency for Healthcare Research and Quality

HCUP NATIONWIDE READMISSIONS DATABASE (NRD) ABSTRACT

The Nationwide Readmissions Database (NRD) is part of the Healthcare Cost and Utilization Project (HCUP) that is sponsored by the Agency for Healthcare Research and Quality (AHRQ). The NRD addresses a large gap in healthcare data – the lack of nationally representative information on hospital readmissions for all patients, regardless of the expected payer for the hospital stay. The NRD was created to enable analyses of national readmission rates and to support public health professionals, administrators, policymakers, and clinicians in their decision making.

The NRD is drawn from HCUP State Inpatient Databases (SID) that contain reliable, verified patient linkage numbers that can be used to track a person across hospitals within a State, while adhering to strict privacy guidelines. The 2019 NRD is constructed from 30 SID. These States are geographically dispersed and account for 61.8 percent of the total U.S. resident population and 60.4 percent of all U.S. hospitalizations. See <u>Appendix A, Table 1</u> for a list of data organizations participating in the NRD.

The NRD includes community hospitals, excluding rehabilitation or long-term acute care hospitals. All discharges from the SID are included except the following:

- Discharges from patients with an age of 0 from 14 of the 30 SID in 2019 because patient linkage numbers are inconsistently reported for this age
- Discharges with missing patient linkage numbers
- Discharges with questionable patient linkage numbers, defined as 20 or more discharges in a year, hospitalized after discharged dead, and overlapping inpatient stays
- Discharges from hospitals with more than 50 percent of their total discharges excluded for any of the above causes, because patients treated at these hospitals may not be reliably tracked over time. This does not include the exclusion for patients at age 0 with verified patient linkage numbers on less than 90 percent of discharges for this age group.

After exclusions, the 2019 NRD contains about 84 percent of SID discharges from the participating States. Unweighted, the NRD contains approximately 18 million discharges in 2019. Weighted, it estimates roughly 35 million discharges in the United States.

The NRD is designed to be flexible to various types of analyses of national readmissions for all payers. The criteria to determine the relationship between multiple hospital admissions for an individual patient are left to the analyst using the NRD. Outcomes of interest include national readmission rates, reasons for returning to the hospital for care, and the hospital costs for discharges with and without readmissions. *The NRD is not designed to support regional, State-, or hospital-specific readmission analyses.*

Access to the NRD is open to users who sign a Nationwide Databases Data Use Agreement

(DUA). Uses are limited to research and aggregate statistical reporting. For more information on the NRD, visit the AHRQ-sponsored HCUP-US website at <u>www.hcup-us.ahrq.gov</u>.

INTRODUCTION TO THE NATIONWIDE READMISSIONS DATABASE (NRD)

Overview of NRD Data

The Healthcare Cost and Utilization Project (HCUP) Nationwide Readmissions Database (NRD) was created to enable analyses of national readmission rates and to support public health professionals, administrators, policymakers, and clinicians in their decision making. Reducing hospital readmissions is a key strategy for improving the quality of healthcare while containing cost. The NRD is designed to be flexible to various types of readmission analyses. The database includes discharges for patients with and without repeat hospital visits in a year and those who have died in the hospital. Repeat visits may or may not be related. The criteria to determine the relationship between hospital admissions are left to the analyst using the NRD.

States Included in the NRD

Some HCUP Partner organizations provided synthetic patient linkage numbers in their SID that can be used to track patients within and across hospitals in a particular State. Unique combinations of the patient linkage number and the patient's date of birth and sex constituted a verified patient linkage number.

States selected for the NRD have verified patient linkage numbers on at least 90 percent of adult discharges. From these States, the NRD included discharges from patients aged 1 year and older. Discharges aged 0 were only retained from States that had verified patient linkage numbers on more than 90 percent of discharges aged 0. <u>Appendix A, Table 1</u> identifies the statewide data organizations that contribute to the NRD. <u>Appendix A, Figure 1</u> displays the geographic distribution of the 30 HCUP Partner organizations participating in the 2019 NRD. Based on U.S. Census Bureau data, the 2019 NRD accounts for 61.8 percent of the U.S. population and 60.4 percent of U.S. hospitalizations reported in the American Hospital Association (AHA) Annual Survey Database.

Hospitals Included in the NRD

The SID contains inpatient discharges from all hospitals provided by the HCUP Partners (e.g., community, general acute care, and specialty hospitals). The American Hospital Association (AHA) defines *community* hospitals as "all nonfederal short-term general and special hospitals, including special children's hospitals, whose facilities and services are available to the public."¹ Specialty hospitals included in the AHA definition of community hospitals are obstetrics-gynecology, ear-nose-throat, short-term rehabilitation, orthopedic, pediatric institutions, and long-term acute care (LTAC) facilities. Also included are public hospitals and academic medical centers.

The NRD is limited to data from community hospitals that are not rehabilitation or LTAC facilities. Noncommunity hospitals were excluded because of inconsistent capture of data across HCUP States. Rehabilitation or LTAC hospitals were excluded because they treat a unique patient population that has longer stays and higher costs. Information on the percentage of SID discharges excluded by type of exclusion is provided in <u>Appendix A, Table 2</u>. Details on the number of hospitals in the NRD are provided in <u>Appendix A, Table 3</u>.

¹ American Hospital Association Glossary (<u>https://www.ahadataviewer.com/glossary/)</u>

Discharges Included in the NRD

All SID discharges from selected States and hospitals were included, with a few exceptions. Records with missing or unverified patient linkage numbers were excluded from the NRD. All discharges aged 0 from States that had verified patient linkage numbers on less than 90 percent of discharges for this age group were excluded.

Another concern was verified patient linkage numbers that did not appear to track an individual within the year for the following reasons:

- (1) Extraordinary utilization in the year, defined as 20 or more admissions in a calendar year
- (2) Multiple discharges for the same identifier that showed the patient discharged dead from one admission and then admitted at a later date in the year
- (3) Overlapping hospitalizations for the same patient linkage number at the same or different hospitals.

Discharges for patient linkage numbers considered questionable because of these three criteria were excluded from the NRD.

Discharge-specific exclusions such as the removal of unverified, missing, or questionable patient linkage numbers impacted individual hospitals if they had more than 50 percent of their annual discharges excluded. These hospitals were not good candidates for a readmission analysis because too many of their discharges could not be tracked over time and were therefore excluded from the NRD. Information on the percentage of SID discharges excluded by type of exclusion is provided in <u>Appendix A, Table 2</u>.

There were no exclusions for certain types of patients or clinical conditions. Therefore, conditions such as childbirth that generally do not result in a readmission were also included. Discharges that resulted in an in-hospital death were retained because these were possible readmission records. Discharges for residents and nonresidents were included in the State in which they were treated. Although most patients seek treatment at hospitals in their State of residence, there are occasions when patients are treated at hospitals in another State. Hospitals that specialize in certain types of care may attract patients from across the United States. In addition, hospitals near State borders frequently treat patients that reside in neighboring States. The data element RESIDENT identifies a discharge as a resident of the State in which he or she received hospital care. Details on the number of discharges in the NRD are provided in <u>Appendix A, Table 3</u>.

Discharges Involving Transfers

Hospital administrative databases like the NRD and SID are "discharge-level" files, meaning that each record represents one discharge abstract from an inpatient stay. If a patient visits the hospital multiple times in a given year, the SID includes separate records for each inpatient stay. In addition, if a patient is transferred between hospitals within the State, the SID contain two discharge records, one record from the first hospitalization and a second record from the latter hospitalization.² Readmission analyses do not usually allow the hospitalization at the receiving hospital to be counted as a readmission. To eliminate this possibility, the pairs of records representing a transfer were collapsed into a single "combined" record in the NRD and removed the original separate discharge records from the NRD. *Transfer* records were defined

² If the patient is transferred to an out-of-state hospital, the subsequent discharge would not be included in the NRD because the HCUP patient linkage numbers only can follow a patient within a State.

as having all of the following characteristics:

- Discharge date of the first inpatient stay equaled the admission date of a subsequent inpatient stay³
- The first record had a discharge disposition of transfer to an acute care hospital
- The second record was from a different hospital and had an admission source indicating a transfer.

There are some records that meet only the first criterion. For this type of record, they were defined as *same-day* stays, that is, the discharge date for one inpatient stay was the same as the admission date of a second stay for the same patient, but there was no indication of a transfer by the discharge disposition or admission source. Same-day stays may or may not have involved different hospitals. Same-day stays may indicate that a patient was discharged too soon and then needed to be returned to the hospital on the same day. However, it was also possible that these were transfer records with an incorrect or missing discharge disposition and admission source. Therefore, the pairs of records representing same-day stays were also collapsed into a single "combined" record. A more detailed description of the methodology used to identify and combine transfer and same-day stays is provided in <u>Appendix B</u>.

Summary of Hospitals and Discharges Included in the NRD

The NRD includes community hospitals and excludes the following types of hospitals:

- Noncommunity hospitals because of inconsistent capture of data across HCUP States
- Community hospitals that are either rehabilitation or LTAC hospitals because these hospitals treat a unique patient population that has longer stays and higher costs.

All discharges from included hospitals were retained in the NRD with the following exceptions:

- Discharges from patients with an age of 0 from SID with verified patient linkage numbers on less than 90 percent of discharges for this age group
- Discharges with missing or unverified patient linkage numbers
- Discharges with questionable patient linkage numbers, defined as 20 or more discharges in a year, hospitalized after discharged dead, and overlapping inpatient stays
- Discharges from hospitals with more than 50 percent of their total discharges excluded for any of the above two causes. This does not include the exclusion for patients with an age of 0 with verified patient linkage numbers on less than 90 percent of discharges for this age group.

Information on the percentage of SID discharges excluded by type of exclusion is provided in <u>Appendix A, Table 2</u>. There were no exclusions for certain types of patients or clinical conditions. Discharges that resulted in an in-hospital death were included because these were possible readmission records. Discharges for residents and nonresidents of the State in which they were treated were also included. In addition, pairs of discharge records representing transfers or same-day stays (i.e., one discharge record from the sending hospital and one

³ Although the text refers to using the discharge and admission date, in reality, the NRD data elements NRD_DaysToEvent and LOS were used to identify the sequential order of inpatient records and whether they stopped and started on the same day.

discharge record from the receiving hospital) were collapsed into a single record so that the hospitalization at the receiving hospital could not be counted as a readmission. Details on the number of States, hospitals, and discharges in the NRD are provided in <u>Appendix A, Table 3</u>.

State-Specific Restrictions

Some sources that contributed data to the NRD imposed restrictions on the release of certain data elements or discharges that could be included in the database. In addition, because of confidentiality laws, some data sources were prohibited from providing HCUP with discharge records that indicated specific medical conditions, such as HIV/AIDS or behavioral health. Detailed information on these State-specific restrictions is available in <u>Appendix C</u>.

ICD-10-CM/PCS Started October 1, 2015 at the Beginning of Fiscal Year 2016

On October 1, 2015, the United States transitioned from using ICD-9-CM to ICD-10-CM/PCS code sets for reporting medical diagnoses and inpatient procedures.⁴ ICD-10-CM/PCS consists of two parts:

- ICD-10-CM: diagnosis coding on inpatient and outpatient data
- ICD-10-PCS: procedure coding on inpatient data.

The HCUP-US website has a section on <u>ICD-10-CM/PCS Resources</u> that summarizes key issues for researchers using HCUP and other administrative databases that include ICD-10-CM/PCS coding. The web page provides general guidance and forewarning to users analyzing outcomes that may be affected by the transition to the ICD-10-CM/PCS coding system and lists other related Web resources.

File Structure of the NRD

Generally, the NRD includes three discharge-level files and one hospital-level file:

Discharge-level files

- Core File is a single file containing commonly used data elements (e.g., age, expected primary payer, discharge status, ICD-10-CM/PCS codes, total charges).
 - This file is available in all years of the NRD.
- **Severity File** is a single file containing additional data elements to aid in identifying the severity of the condition for a specific discharge.
 - This file is available in all years of the NRD.
- Diagnosis and Procedure Groups File is a single file containing additional information on the ICD-10-CM diagnoses and ICD-10-PCS procedures created by the HCUP software tools.
 - For data years 2016–2017, this file was not available in the NRD. Data elements derived from the ICD-10-CM/PCS HCUP software tools are not included in the NRD because they were still in development and testing.

⁴ ICD-9-CM: International Classification of Diseases, Ninth Revision, Clinical Modification;

ICD-10-CM/PCS: International Classification of Diseases, Tenth Revision, Clinical Modification/Procedure Coding System

- Beginning with data year 2019, this file is available in the NRD and includes data elements derived from the Clinical Classifications Software Refined (CCSR) for ICD-10-CM diagnoses.
- Beginning with data year 2019, data elements derived from the Elixhauser Comorbidity Software Refined for ICD-10-CM, the CCSR for ICD-10-PCS procedures, and Procedure Classes Refined for ICD-10-CM are also available in this file.

Hospital-level file

- **Hospital File** is a single file containing information on hospital characteristics.
 - This file is available in all years of the NRD.

The NRD unique record identifier (HCUP data element KEY_NRD) can be used to add data elements from the Severity file and Diagnosis and Procedure Groups file, if available, to the records on the Core file. The NRD hospital identifier (HCUP data element HOSP_NRD) can be used to add data elements from the hospital-level file to the Core file.

File Structure of the 2015 NRD

The NRD data files are annual, calendar-year files based on discharge date for all years except 2015. The introduction of ICD-10-CM/PCS on October 1, 2015 means that the 2015 NRD includes a combination of codes:

- Nine months of the data with ICD-9-CM codes (January 1, 2015 to September 30, 2015)
- Three months of data with ICD-10-CM/PCS codes (October 1, 2015 to December 31, 2015).

To alert users to this change in the ICD coding scheme, the file structure of the 2015 NRD differs from the annual files for other data years in three primary ways:

- The names of diagnosis- and procedure-related data elements under ICD-10-CM/PCS have been changed to identify the coding scheme with a prefix of "I10_."
- Diagnoses and procedures, and related data elements, have been moved out of the Core File and into the Diagnosis and Procedure Groups File where the first three quarters of data (with ICD-9-CM codes) are stored separately from the fourth quarter of data (with ICD-10-CM/PCS codes).
- Data elements based on the HCUP software tools that are derived from ICD-10-CM/PCS codes are not included in the fourth quarter data.

More information about the file structure of the 2015 NRD is available in the <u>Introduction to the</u> <u>Nationwide Readmissions Database (NRD), 2010–2015</u> and on the <u>NRD Database</u> <u>Documentation</u> page of the HCUP-US website.

NRD Data Elements

The coding of data elements in the NRD is consistent with other HCUP databases. The following three objectives guided the definition of data elements in all HCUP databases:

• Ensure usability without extensive editing by analysts.

- Retain the largest amount of information available from the original sources, while still maintaining consistency among sources.
- Structure the information for efficient storage, manipulation, and analysis.

More information on the coding of HCUP data elements is available on the HCUP User Support (HCUP-US) website (<u>www.hcup-us.ahrq.gov/db/coding.jsp</u>).

The NRD contains more than 100 clinical and non-clinical data elements provided in a hospital discharge abstract, such as:

- Discharge month, quarter, and year⁵
- ICD-10-CM diagnoses (or ICD-9-CM for discharges prior to October 1, 2015)
- ICD-10-PCS procedures (or ICD-9-CM for discharges prior to October 1, 2015)
- Patient demographics (e.g., sex, age, income quartile based on the median household income of the patient's ZIP Code, and urban/rural location of the patient's residence)
- Expected source of primary payment (e.g., Medicare, Medicaid, private insurance, selfpay, and other insurance types)
- Total hospital charges and hospital cost⁶
- Discharge weights for generating national estimates.
- Data elements essential to readmission analyses:
 - Verified patient linkage number (HCUP data element NRD_VisitLink) that can be used to identify discharges belonging to the same patient
 - Timing between admissions for a patient (HCUP data element NRD_DaysToEvent)
 - Length of inpatient stay in days (HCUP data element LOS)
 - Identification of transfers, same-day stays, and combined transfer records (HCUP data elements SAMEDAYEVENT and REHABTRANSFER)
 - Identification of the patient as a resident of the State in which he or she received hospital care (HCUP data element RESIDENT).
- Data elements derived from the HCUP software tools for ICD-10-CM/PCS beginning data year 2018 or the HCUP software tools for ICD-9-CM through quarter 3 of data year 2015.⁷

<u>Appendix D</u> identifies the data elements in each 2019 NRD file:

• <u>Table 1</u> for the NRD Core File

⁵ Admission and discharge dates were not available because of patient confidentiality restrictions.

⁶ Total hospital cost must be added to the NRD using the HCUP supplemental Cost-to-Charge Ratio files.

⁷ Users interested in applying HCUP software tools to the NRD to produce data elements unavailable in the database files for a certain year may do so by downloading the respective tool(s) from the <u>Research</u> <u>Tools</u> section of the HCUP User Support (HCUP-US) website. Additionally, users may wish to review the <u>HCUP Software Tools Tutorial</u>, which provides instructions on how to apply the HCUP software tools to HCUP or other administrative databases.

- <u>Table 2</u> for the NRD Severity Files
- <u>Table 3</u> for the NRD Diagnosis and Procedure Groups Files
- <u>Table 4</u> for the Hospital File.

The tables in <u>Appendix D</u> provide summary documentation for the data. Please refer to the NRD documentation located on the HCUP-US website (<u>www.hcup-us.ahrq.gov</u>) for comprehensive information about data elements and the files.

For prior years, refer to the <u>NRD Description of Data Elements</u> page on the HCUP-US website or to previous versions of the NRD Introduction.

Getting Started

The HCUP Nationwide Readmissions Database (NRD) is distributed as comma-separated value (CSV) files delivered via secure digital download from the <u>Online HCUP Central Distributor</u>. The files are compressed and encrypted with SecureZIP® from PKWARE.

The NRD product is downloaded in a single zipped file for each year which contains several data-related files and accompanying documentation. The data-related files include the following compressed files:

- 1) Core File (NRD_yyyy_Core.zip, where yyyy indicates the data year)
- 2) Severity Measures File (NRDS_yyyy_Severity.zip, where yyyy indicates the data year).
- 3) Diagnosis and Procedure Groups File (NRD_yyyy_DX_PR_GRPS.zip, where yyyy indicates the data year)
- 4) Hospital File (NRD_yyyy_Hospital.zip, where yyyy indicates the data year)

To load and analyze the NRD data on a computer, users will need the following:

- The password provided by the HCUP Central Distributor
- A hard drive with at least 71 gigabytes of space available
- A third-party zip utility such as ZIP Reader, SecureZIP®, WinZip®, or Stuffit Expander®
- SAS®, SPSS®, Stata® or similar analysis software.

The total size of the comma-delimited version of the 2019 NRD is 41 GB. The 2019 NRD files loaded into SAS are about 10 GB with compression. In SAS, the largest use of space typically occurs during PROC SORT, which requires workspace about three times the size of the file. Thus, the NRD files would require at least 30 GB of available workspace to perform a sort procedure. Most SAS data steps will require twice the storage of the file, so that both the input and output files can coexist. The NRD files loaded into SPSS are under 150 GB. Because Stata loads the entire file into memory, it may not be possible to load every data element in the NRD Core file into Stata. Stata users will need to maximize memory and use the "_skip" option to select a subset of data elements. More details are provided in the Stata load programs.

With a file of this size and without careful planning, space could easily become a problem in a multistep process. It is common to produce several versions of a file during data preparation, as well as further multiple versions for analysis. Therefore, the amount of space required could escalate rapidly.

Decompressing the NRD Files

To extract the data files from the compressed download file, follow these steps:

- 1) Create a directory for the NRD on your hard drive.
- 2) Unzip the compressed NRD product file into the new directory using a third-party zip utility. This will place compressed, encrypted data-related files in the new directory. You will be prompted to enter the encryption password (sent separately by email) to decrypt the file.

Please note that attempts to unzip encrypted files using the built-in zip utility in Windows® (Windows Explorer) or Macintosh® (Archive Utility) will produce an error message warning of incorrect password and/or file or folder errors. The solution is to use a third-party zip utility.

Third-party zip utilities are available from the following reputable vendors on their official Web sites.

- ZIP Reader (Windows) (free download offered by the PKWARE corporation)
- SecureZIP® for Mac or Windows (free evaluation and licensed/fee software offered by the PKWARE corporation)
- WinZip (Windows) (evaluation and fee versions offered by the WinZip corporation)
- Stuffit Expander® (Mac) (free evaluation and licensed/fee software offered by Smith Micro corporation)
- Unzip each of the compressed, encrypted data-related files using the same password and third-party zip utility method. This will place the data-related ASCII files in this same directory by default.

Downloading and Running the Load Programs

Programs to load the data into SAS, SPSS, or Stata, are available on the HCUP User Support website (HCUP-US). To download and run the load programs, follow these steps:

- 1) Go to the NRD Database Documentation page on HCUP-US at <u>www.hcup-us.ahrq.gov/db/nation/nrd/nrddbdocumentation.jsp</u>.
- 2) Go to the "File Specifications and Load Programs" section on this page.
- 3) Click on "Nationwide SAS Load Programs", "Nationwide SPSS Load Programs", or "Nationwide Stata Load Programs" to go to the corresponding Load Programs page.
- Select the data year and the database ("NRD") from the drop-down lists on this page. Or you may select "NRD Load All Years" to obtain a zipped file with all load programs for multiple years at once.
- 5) Select and save the load programs you need. The load programs are specific to the data year and data-related file. For example, the load program for the 2019 NRD Core file is found under the link "SAS NRD 2019 Core File" in the list generated by selecting "2019" and "NRD." Save the load programs into the same directory as the NRD CSV files on your computer.

6) Edit and run the load programs as appropriate for your computing environment to create the analysis files. For example, modify the directory paths to point to the location of your input and output files.

NRD Documentation

Comprehensive documentation for the NRD files is available on the HCUP-US website (<u>www.hcup-us.ahrq.gov/db/nation/nrd/nrddbdocumentation.jsp</u>). Users of the NRD can access complete file documentation, including variable notes, file layouts, load programs, and summary statistics. Refer to these resources to understand the structure and content of the NRD and to aid in using the database.

Appendix A, Table 4 details the comprehensive NRD documentation available on HCUP-US.

HCUP Online Tutorials

For additional assistance, AHRQ has created the HCUP Online Tutorial Series, a series of free, interactive courses that provide training on technical methods for conducting research with HCUP data. Topics include an <u>HCUP Overview Course</u> and these tutorials that are helpful to NRD data users:

- The <u>Nationwide Readmissions Database Tutorial</u> provides a description of the NRD database design and a detailed example on how to use the NRD to estimate national readmission rates with standard errors.
- The <u>Load and Check HCUP Data</u> tutorial provides instructions on how to unzip (decompress) HCUP data, save it on a computer, and load the data into a standard statistical software package. This tutorial also describes how to verify that the data have been loaded correctly.
- The <u>HCUP Software Tools Tutorial</u> introduces users to the HCUP software tools, which can be applied to HCUP and other administrative databases to create new data elements from existing data, thereby enhancing a researcher's ability to conduct analyses. There are four modules within this course grouping the HCUP tools by the following coding systems: ICD-10-CM diagnoses, ICD-10-PCS procedures, Current Procedural Terminology (CPT[®]) and Healthcare Common Procedure Coding System (HCPCS) Level II codes, and ICD-9-CM diagnoses and procedures. Users will learn about the purpose of each tool and receive technical guidance for applying the tools to their data.

Other tutorials about the design or use of other HCUP databases also are available. The Online Tutorial Series is located on the HCUP-US website at <u>www.hcup-us.ahrq.gov/tech_assist/tutorials.jsp</u>.

HCUP Reports on Readmission Methodology

Several HCUP Statistical Briefs have been published that provide statistics about national readmission rates, including readmissions involving specific diagnoses and procedures. A full list of these available reports is available in <u>HCUP Statistical Briefs by Topic</u> – Readmissions and Revisits.

Three HCUP Methods Series reports have been released that provide additional information on readmissions:

- Overview of Key Readmission Measures and Methods (<u>Report #2012-04</u>). This report describes 12 key measures of hospital readmissions from different agencies or companies. The 12 measures were selected for inclusion because they are: (1) endorsed by the National Quality Forum (NQF), (2) currently used by AHRQ in various initiatives, or (3) otherwise well-known and used by researchers.
- Methodological Issues when Studying Readmissions and Revisits using Hospital Administrative Data (Report #2011-01). This report discusses the decisions that need to be addressed as an analyst designs a study of patients with sequential acute care hospital visits based on hospital administrative data. Topics include considerations in preparing the administrative database for the analysis, defining repeat hospital visits, and reporting results.
- Calculating Nationwide Readmission Database (NRD) Variances (<u>Report #2017-01</u>). This report provides guidance on calculating variances for estimates of readmission outcomes using the NRD.

HCUP has also created a topical report that provides insight on how the ICD-10-CM/PCS transition has affected readmission rates:

 Impact of the ICD-10-CM/PCS Transition on Readmission Rates, 2015-2017. The report describes the methodology and findings of an analysis that examined the impact of the transition from the ICD-9-CM to the ICD-10-CM/PCS coding system on condition-specific readmission statistics. Conditions are based on two categorizations—DRGs and major diagnostic categories (MDCs).

ICD-9-CM and ICD-10-CM/PCS Diagnosis and Procedure Codes

The NRD includes ICD-9-CM diagnosis and procedure codes on inpatient discharges prior to October 1, 2015. Starting on October 1, 2015, diagnosis and procedure codes are reported using ICD-10-CM/PCS. HCUP has developed recommendations for reporting statistics (e.g., counts, rates, averages) that are based on HCUP data with a mixture of ICD-9-CM and ICD-10-CM/PCS codes. These recommendations apply to calendar year 2015 data (which includes both ICD-9-CM and ICD-10-CM/PCS codes), as well as reporting trends that span the October 1, 2015 transition date (before and after the introduction of ICD10-CM/PCS).

ICD-9-CM and ICD-10-CM diagnosis and ICD-10-PCS procedure codes provide valuable insights into the reasons for hospitalization and what procedures patients receive, but these codes need to be carefully used and interpreted. ICD-9-CM and ICD-10-CM/PCS codes change every October as new codes are introduced, and some codes are retired. It is critical to check all ICD-9-CM and/or ICD-10-CM/PCS codes used for your analysis to ensure the codes are in effect during the time period studied.

SAMPLING DESIGN OF THE NRD

The NRD was built to facilitate analyses of both all-cause and condition-specific readmissions. National estimates can be produced by applying weighting and stratification methods.

Target Universe

The target universe was limited to inpatient discharges treated at community hospitals in the United States that were not rehabilitation or LTAC facilities. Information on the target universe of these hospitals was available from the American Hospital Association (AHA) Annual Survey of Hospitals. The AHA Survey includes information on the number of inpatient admissions plus births and hospital characteristics such as ownership, number of beds, and location.

Sampling Frame

The sampling frame for the NRD was limited to discharges for patients treated at community hospitals in the NRD States that were not rehabilitation or LTAC facilities. All of the discharges in the sampling frame were included, making the NRD a sample of convenience. Sampling discharges or hospitals was not recommended because the sample needed to balance the database's ability to estimate readmissions for common conditions such as chronic illnesses with the ability to estimate readmissions for rare diseases such as sickle cell anemia. Developing the database using a 100 percent sample allows researchers to study both all-cause and condition-specific readmissions.

Discharge Weights

This section explains the need for post-stratification for weighting the sampling frame to the target universe and the weighting strategy. The term *post-stratification* is used because the stratification was performed after the data exclusions. Discharge weights for national estimates were developed using the target universe as the standard. Although discharge-level weights were calculated for the NRD, hospital-level weights were not calculated. The NRD is not designed to support hospital-specific analyses. The NRD is a 100 percent sample of discharges, not hospital; hospital weights are not applicable.

Post-Stratification for Weighting

Post-stratification for the purpose of weighting allows for compensation of any over- or underrepresented types of hospitals and discharges in the sampling frame (the NRD) with respect to the distribution in the target universe (AHA data). The NRD was post-stratified by hospital and patient characteristics. Consistent with the National Inpatient Sample (NIS) design, the following hospital characteristics explained significant differences in inpatient outcomes: census region, urban/rural location, hospital teaching status, size of the hospital defined by the number of beds, and hospital ownership.⁸

Discharges with unverified and missing patient linkage numbers were excluded because these patients cannot be tracked across time. In an examination of the distribution of patient age and sex between discharges with verified and unverified/missing patient linkage numbers in data year 2011, it was determined that the majority of the unverified discharges (52.7 percent) were females aged 18–44 years old. In addition, 30-day all-cause readmission rates in 2011 by age–sex categories were compared across States, which revealed variation between SID with a high percentage of verified patient linkage numbers and States with a lower percentage of verified patient linkage numbers. These analyses demonstrated that there were differences between

⁸ Changes in the NIS Sampling and Weighting Strategy for 1998. ONLINE January 18, 2002. Available at <u>www.hcup-us.ahrq.gov/db/nation/nis/reports/Changes_in_NIS_Design_1998.pdf</u>. Accessed September 15, 2011.

discharges with and without patient linkage numbers by sex and age. Therefore, the NRD was also post-stratified by sex and five age groups (0, 1–17, 18–44, 45–64, and 65 and older).

Weighting

The target universe totals for the discharge counts by stratum were based on all SID discharges from all HCUP Partners, unless there were missing hospitals. If there were hospitals missing from the stratum according to the AHA, then the target universe total included SID discharges for all available hospitals plus the AHA admission counts for the missing hospitals. This approach was consistent with the NIS and took advantage of the fact that the SID included over 95 percent of discharges from community hospitals that are not rehabilitation or LTAC hospitals in the United States. Discharge counts for the sampling frame were based on the NRD discharges (after the exclusion of States, hospitals, and discharges).

To determine discharge-level weights, the number of discharges for the target universe and the sampling frame were summarized by stratum. Each stratum was defined by hospital characteristics (census region, urban/rural location, hospital teaching status, size of the hospital defined by the number of beds, and type of ownership) and patient characteristics (sex and five age groups [0, 1–17, 18–44, 45–64, and 65 and older]). Within each stratum, *s*, each NRD inpatient admission received a weight:

DISCWT_{i,j} = Ns(universe) _{i,j} ÷ Ns(sample)_{i,j}

where Ns(universe) $_{i,j}$ represents the number of inpatient discharges at community hospitals that were not a rehabilitation or LTAC hospital in the universe within stratum *s* for sex *i* and age group *j*; Ns(sample) $_{i,j}$ is the number of inpatient discharges in the sampling frame for sex *i* and age group j. Age group *j* included ages 0, 1–17, 18–44, 45–64, 65 and older. Therefore, each discharge's weight (DISCWT_{*i*,*j*}) is equal to the number of inpatient discharges it represents in stratum *s* for sex *i* and age group *j* during that year.

To improve reliability of the age distribution of the SID discharges, some strata in the target universe were collapsed prior to the weight calculations such that at least two SID hospitals and at least 100 discharges from the SID were included in each stratum. In addition, some strata were collapsed to include at least two hospitals in each stratum in the sample frame. Strata were first collapsed across control/ownership, combining either the two private designations or all three types of control (public, private not-for-profit, and private for-profit). If the stratum combined across control still lacked a sufficient number of hospitals or discharges, then the location category was collapsed. Small and large metropolitan areas were combined or micropolitan and rural areas were combined. Lastly, if the stratum still lacked a sufficient number of hospitals or discharges, then the bed-size category was collapsed with large and medium hospitals combined. There was no collapsing of strata across region or teaching status. In addition, weights were adjusted if a hospital was missing data for one or more quarters in the year. The range of weights by age and sex are provided in <u>Appendix A, Table 5</u>.

Final NRD Sample

In summary, the NRD is an annual file constructed using one calendar year of discharge data. Included discharges were treated at community hospitals (excluding rehabilitation or LTAC hospitals) for which the majority of their discharges had patient linkage numbers that were verified and not questionable. Discharge weights were calculated using post-stratification on hospital characteristics (census region, urban-rural location, teaching status, bed size, and hospital control) and patient characteristics (sex and five age groups [0, 1–17, 18–44, 45–64, and 65 and older]). The target universe of inpatient discharges in the United States was estimated for each stratum using SID total discharges augmented by AHA discharge counts when hospitals were not reported in the SID. Details on the number of States, hospitals, and discharges in the NRD are provided in <u>Appendix A, Table 3</u>. The range of discharge weights by age and sex are provided in <u>Appendix A, Table 5</u>. The NRD is designed to be flexible to various types of analyses of national readmissions. *The NRD is not designed to support regional, State- or hospital-specific readmission analyses*.

Limitations of the NRD

The NRD contains about 18 million inpatient discharge records and over 100 clinical and nonclinical data elements. A multitude of research studies can be conducted with the data, but there are some limitations.

- Limitations on Studying Pediatric Readmissions: The NRD excludes discharges for patients who were younger than 1 year from two-thirds of the SID in 2010–2012 NRD, about 60 percent of the SID in the 2013 NRD, and about half of the SID in the 2014-2019 NRD. The weights for pediatric discharges were often higher than adult discharges. The range of discharge weights by age and sex are provided for each data year in <u>Appendix A, Table 5</u>.
- Limitations from Using One Year of Discharge Data: The NRD is an annual file containing inpatient records for patients discharged in a calendar year. The files included patients admitted in the prior year and discharged in the current year, while excluding patients admitted to a hospital in the current year but discharged in the next year. Therefore, 30- or 60-day readmissions for patients admitted in the latter part of the year may not be captured if the subsequent admission crossed into the next year. In addition, one year of discharge data is probably an insufficient length of time for examining readmissions that are more than 90 days apart.
- Limitations about Using More than One Year of the NRD: The 2010–2019 NRD cannot be combined across data years to create a multi-year database. The patient linkage numbers (NRD_VisitLink) do not track the same person across years. The hospital identifiers in HOSP_NRD do not track the same hospital across years. Each year of the NRD must be considered as a separate sample.
- Limitations from Using State-Specific Identifiers: Patients who were hospitalized in one State and readmitted or transferred to a hospital in another State cannot be tracked in the NRD, because each of the SID uses different coding for their patient linkage numbers. The NRD includes nonresident patients because discharges at border hospitals that provided care for patients in their community were retained, even though that community happened to cross State borders. To understand the impact on readmission rates caused by using State-specific patient linkage numbers, a sensitivity analysis were performed using the 2011 Medicare Standard Analytic File (SAF) that allowed admissions for patients covered by Medicare fee-for-service (FFS) plans to be tracked across all States.

The analysis demonstrated that condition-specific readmission rates were higher if a patient could be tracked across all States, but the percentage increase was less than 5 percent for most of the CCS categories. This analysis was limited to the Medicare FFS

population because of the data source. The Medicare (FFS and managed care) population accounted for about 40 percent of all inpatient discharges in 2011,⁹ and previous research indicates that the Medicare population has higher readmission rates than populations covered by for other payers.¹⁰ Conditions often associated with younger adults, such as pregnancy, were included in the Medicare estimates because about 20 percent of Medicare discharges are younger than 65 years.¹¹ Medicare patients younger than 65 include people who are disabled or who have been diagnosed with end-stage renal disease or amyotrophic lateral sclerosis (ALS).

Given the volume and severity of illness for Medicare patients, the estimates for the increase in the condition-specific readmission rates using the Medicare data provide a reasonable upper boundary on the impact. The details of this analysis are provided in Appendix E.

HOW TO USE THE NRD FOR READMISSION ANALYSES

This section provides a brief synopsis of special considerations for using the NRD. For more details, refer to the comprehensive documentation on the HCUP-US website (<u>www.hcup-us.ahrq.gov/db/nation/nrd/nrddbdocumentation.jsp</u>).

All persons using the NRD (whether or not they are the original recipient of the data) must complete the online Data Use Agreement Training Tool available on the HCUP-US Web site (<u>www.hcup-us.ahrq.gov/tech_assist/dua.jsp</u>) and then read and sign a Data Use Agreement. A copy of the signed Data Use Agreements must be sent to the HCUP Central Distributor. See page two of this document for the mailing address.

Select readmission statistics are included in Appendix A, Tables 6-8.

⁹ HCUPnet query on the expected primary payer for the 2011 Nationwide Inpatient Sample. Accessed December 9, 2014.

¹⁰ Wier LM, Barrett ML, Steiner C, Jiang HJ. All-Cause Readmissions by Payer and Age, 2008. HCUP Statistical Brief No. 115. June 2011. Rockville, MD: Agency for Healthcare Research and Quality. www.hcup-us.ahrq.gov/reports/statbriefs/sb115.pdf.

¹¹ HCUPnet query on the expected primary payer and age for the 2011 Nationwide Inpatient Sample. Accessed December 9, 2014.

NRD Data Elements Critical to Tracking a Patient and Determining the Time Between Admissions

For any readmission analysis of inpatient stays, three HCUP data elements are critical to tracking a patient and determining the time between admissions: NRD_VisitLink, NRD_DaysToEvent, and LOS (length of stay).

Patient Linkage Number (NRD_VisitLink)

NRD_VisitLink is the data element that links for all inpatient stays associated with a unique patient. All discharges in the NRD include a value for NRD_VisitLink. The value was assigned based on a unique combination of the synthetic patient linkage number provided by the HCUP Partner, date of birth, and sex. No verified patient linkage number was assigned if any one of the three pieces of information was missing. Because of discharge-level exclusions, NRD_VisitLink is always coded on records in the NRD.

Although the term *patient linkage number* is used to describe the information in the NRD data element NRD_VisitLink, the values are not recognizable as specific patient information. NRD_VisitLink does not include the values of the encrypted person's social security number, date of birth, or sex.

Time Between Admissions (NRD_DaysToEvent and LOS)

NRD_DaysToEvent is the number of days from a randomly chosen "start date" to the admission date for each patient's discharge. The actual admission and discharge dates could not be included on the NRD because they were considered highly sensitive information according to Health Insurance Portability and Accountability Act (HIPAA) guidelines. The coding scheme for NRD_DaysToEvent was designed to adhere to these strict privacy guidelines and protect patient confidentiality.

Each verified patient linkage number (NRD_VisitLink) was assigned a unique randomly chosen start date that was used to calculate NRD_DaysToEvent for all visits associated with that NRD_VisitLink value. The data element NRD_DaysToEvent was the difference between the visit's admission date and the start date associated with the NRD_VisitLink. NRD_DaysToEvent was reported as missing if the admission date was unavailable.

For readmission analyses, determining the number of days between the end of one admission and the start of the next admission is critical. There is no data element specific to this timing difference in the NRD because the calculation is dependent on which two discharges are of interest for the readmission study. For example, a study of readmissions for diabetes might only consider the number of days between two diabetes discharges, whereas a study of post-surgery infections might consider any discharge in 30 days.

Because NRD_DaysToEvent was based on the admission date, the calculation of days needs to be the difference of NRD_DaysToEvent between two selected discharges for a unique verified patient linkage number (NRD_VisitLink), adjusted for the length of stay. Consider the following example:

• A patient with congestive heart failure has a 3-day hospital admission on 1/10/2013 and another admission on 1/25/2013.

- The NRD_DaysToEvent value is 1109 for the 1/10/2013 admission, and the NRD_DaysToEvent value is 1124 for the 1/25/2013 admission.
- The number of days between the <u>start of each admission</u> is 15 days (1124 1109 = 15) because NRD_DaysToEvent is based on the admission date.
- The number of days <u>between the admissions</u> (from discharge date of the first admission to the start of the second admission) is 12 days (1124 1109 3 = 12) because the patient had a 3-day LOS.

If NRD_DaysToEvent or LOS was missing, then it was not possible to determine the number of days to a subsequent admission. However, this rarely occurs. LOS was only critical if it was missing on the first admission in a series. If the admission was the second in the series, then LOS was not pertinent.

For a single patient with multiple inpatient stays, the lowest value of NRD_DaysToEvent corresponds to the earliest inpatient stay in the year. It is important to remember that if patient A has a value of 605 for NRD_DaysToEvent and patient B has a value of 300 for NRD_DaysToEvent, patient B's hospital stay did not necessarily take place prior to patient A's stay. In fact, patient B's NRD_DaysToEvent value has no relation to patient A's NRD_DaysToEvent value. Because of the use of a random start date in the calculation of NRD_DaysToEvent, the value of NRD_DaysToEvent cannot be compared across patients.

Additional information about the HCUP revisit variables is available on the HCUP User Support Web site (<u>www.hcup-us.ahrq.gov/toolssoftware/revisit/revisit.jsp</u>).

Defining Readmissions

The NRD was designed to support many different types of readmission analyses. Analysts can use the information contained in the NRD to define the index event and readmission specific to their topic of interest. Common terminology is first defined:

- Index event the starting point for analyzing repeat hospital visits
- *Readmission* a subsequent inpatient admission within a specified time period; readmission may be for a specific cause or any cause.

The next section discusses the following analytic considerations for defining index events and readmissions:

- Defining the index event
- Specifying the criteria for a readmission
- Selecting the appropriate time period to qualify the readmission
- Reporting readmission rates.

Defining the Index Event

The index event is typically defined by a combination of clinical and demographic criteria. Inclusion and exclusion criteria should be used to define an index event indicator that identifies NRD discharges as an index event specific to the analysis of interest. The NRD does not include a data element for index events because they are specific to each analysis. However, the NRD does include the information necessary to define different types of index events. Criteria can include, but are not limited to, age of the patient and specific diagnoses and/or procedures. The NRD contains various data elements that can be used for inclusion criteria:

- ICD-10-CM diagnosis and ICD-10-PCS procedure codes (or ICD-9-CM for discharges prior to October 1, 2015)
- CCSR for ICD-10-CM diagnosis or ICD-10-PCS procedure categories (or CCS for ICD-9-CM categories prior to October 1, 2015)
- Diagnosis-related group (DRG) and major diagnostic category (MDC)
- 3M All-Patient Refined DRG.

Possible exclusion criteria include the following:

- Index events in which the patient died in the hospital, because there is no risk of readmission.
- Patients with complicating comorbidities such as cancer or an immunocompromised state, because these conditions would greatly increase the risk of readmission.
- Transfers and same-day stays that were combined, because they represent a more complex type of care (NRD data element SAMEDAYEVENT not equal to 0).
- Discharge month for when the index event occurred, because there are limitations when using a calendar-year file defined by discharge date.

The annual NRD files include inpatient stays that were discharged in that data year. For example, the 2016 NRD includes admissions that began in 2015 and were discharged in 2016. In contrast, admissions that began later in 2016 and extended into 2017 are not included in the 2016 NRD. About one percent of discharges in the NRD started in the previous year; therefore, it was expected to miss about one percent of admissions that started in the data year and were subsequently discharged in the following year. This will be true for each year of the NRD.

Deciding which discharge months should be included when qualifying an index event depends on (1) the time that will be allowed for a readmission, and (2) the data year. For example, if studying 30-day readmissions with the 2010 - 2014 and 2016 and future years of the NRD, the index event might be limited to those occurring in the discharge months of January through November. That allows the month of December for 30 days of follow-up.¹² In contrast, defining index events using the 2015 NRD needs to take into consideration the transition to ICD-10-CM/PCS on October 1, 2015. For example, if inclusion criteria for index events are based on ICD-9-CM diagnosis or procedure codes, these codes will only be available for the first nine months of the year. As such, it may be advisable to limit diagnosis- or procedure-specific readmissions analyses to these nine months. Selection of the time period will depend on whether the study is focused on diagnosis- and/or procedure-specific index events and readmissions. Consider three examples:

- Example 1. Index events are not defined by diagnoses and/or procedure codes followed by all-cause 30-day readmissions
 - Index events <u>not</u> defined by ICD-9-CM codes but other criteria (e.g., expected payer, age): Limit the index event to the discharge months of January through November. This allows December to identify all-cause 30-day readmissions.

¹² Although it would be advantageous to be able to select a more specific date for a cut-off, patient confidentially concerns limited the available information on the admission and discharge dates to discharge month (data element DMONTH) and discharge year (data element YEAR).

- All-cause readmissions: Use discharge months of January through December. Twelve months can be included because all-cause readmissions are not dependent on diagnosis/procedure coding.
- Example 2. Index events defined by diagnoses and/or procedure codes followed by allcause 30-day readmissions
 - Index events defined by ICD-9-CM codes (e.g., diabetes, hip replacement): Limit the index event to the discharge months of January through September with ICD-9-CM data.
 - All-cause readmissions: Use discharge months of January through October. October data with ICD-10-CM/PCS data can be included because all-cause readmissions are not dependent on diagnosis/procedure coding.
- Example 3. Both index events and 30-day readmissions are defined by diagnoses and/or procedure codes
 - Index events defined by ICD-9-CM codes (e.g., diabetes, hip replacement): Limit the index event to the discharge months of January through August with ICD-9-CM data.
 - Readmissions based on ICD-9-CM codes (e.g., post-surgical infection): Use discharge months of January through September (with ICD-9-CM codes).
 October through December should be excluded from consideration unless the ICD-10-CM/PCS codes can be identified that are equivalent to the ICD-9-CM codes used to define readmissions in January through September.

Specifying the Criteria for a Readmission

Readmission analyses tend to consider one of the following: any subsequent admission regardless of cause, any subsequent admission that does not involve trauma, or any subsequent admission only if the event is "related" to the index event. In addition, a study may consider all readmissions within a time period or just the first readmission. The selection of criteria can dramatically change results. More information on how the results can change is available in an HCUP Method Series report on *Methodological Issues when Studying Readmissions and Revisits Using Hospital Administrative Data*.¹³

The NRD includes several different diagnosis and procedure-related data elements that can be used to examine why a patient returned for hospital care. **The NRD does not identify any discharge as a readmission;** instead, the information necessary to select the appropriate readmission discharges is included in the NRD. Inclusion and exclusion criteria should be used to define a readmission indicator that identifies NRD discharges as readmissions specific to the analysis of interest.

Selecting the Time Period for Revisits

When determining an appropriate time period for the readmission, considerations include selecting a time that encompasses the same risk of exposure to all patients, seasonality of the

¹³ Barrett M, Steiner C, Andrews R, Kassed C, Nagamine M. Methodological Issues when Studying Readmissions and Revisits Using Hospital Administrative Data. HCUP Methods Series Report No. 2011-01. Online March 9, 2011. Rockville, MD: U.S. Agency for Healthcare Research and Quality. <u>www.hcup-us.ahrq.gov/reports/methods/2011_01.pdf</u>.

disease, and possible external factors. Shorter time frames (7 or 14 days) are often used to make events attributable to hospital acute care; longer time frames may reflect differences in ambulatory care and/or coordination of care.

Reporting Rates of Readmission

Although the definition of *readmission rate*—number of readmissions divided by number of cases followed— seems simple, our research into readmission rates showed no standard definition. In some cases, the unit of observation was a patient; in others, the unit of observation was an index event, and individual patients were counted more than once. Some studies focused on the first readmission following an index event, whereas others counted all readmissions. The definitions of the readmission rate were specific to the purpose of the analyses.

Severity or risk adjustment may also be beneficial when comparing readmission rates across geographical regions, hospital types, or different patient populations. A simple risk adjustment would include the age and sex of the patient. A more complex adjustment might also include comorbidites, severity classified by the 3M All-Patient Refined DRG severity score, patient income quartile, or any other factor that could considerably increase or decrease the risk of subsequent hospital care. The NRD includes data elements to support these types of analyses in the Core, Severity, and Diagnosis and Procedure Groups files, if available.

Calculating Nationally Weighted Estimates

An analyst using the NRD must use the discharge-level weight (DISCWT) to produce national estimates. Weighted statistics estimate discharges treated at community hospitals (excluding rehabilitation and LTAC facilities) in the United States. Users should not equate the discharge-level weights to weights that can be used to estimate population counts.

Variance Calculations

It may be important for researchers to calculate a measure of precision for national estimates based on the NRD. Variance estimates must take into account both the sampling design and the form of the statistic. It is important to understand that the NRD is a sample of convenience from the SID and not a sample of hospitals or discharges. Standard error calculations should take into account the stratified sample (data element NRD_STRATUM) and hospitals defining the clusters (data element HOSP_NRD). One resource for understanding the issues surrounding variance calculations for the NRD is the HCUP Method Series Report on *Calculating Nationwide Readmissions Database (NRD) Variances.*¹⁴

To accurately calculate variances from the NRD, appropriate statistical software and techniques must be used. A multitude of statistics can be estimated from the NRD data. Several computer programs that calculate statistics and their variances from sample survey data <u>are listed in the next section</u>. Some of these programs use general methods of variance calculations (e.g., the jackknife and balanced half-sample replications) that take into account the sampling design. However, it may be desirable to calculate variances using formulas specifically developed for certain statistics.

¹⁴ Yoon F, Sheng M, Jiang HJ, Steiner CA, Barrett ML. Calculating Nationwide Readmissions Database (NRD) Variances. HCUP Methods Series Report # 2017-01. Online January 24, 2017. U.S. Agency for Healthcare Research and Quality. <u>www.hcup-us.ahrq.gov/reports/methods/methods.jsp</u>.

Computer Software for Weighted and Variance Calculations

The NRD discharge weights are needed to calculate national estimates of readmission counts and rates. In most cases, computer programs are readily available to perform these calculations. Several statistical programming packages allow weighted analyses.¹⁵ For example, nearly all SAS procedures incorporate weights. In addition, several statistical analysis programs have been developed to specifically calculate statistics and their standard errors from survey data. Version 8 or later of SAS contains procedures (PROC SURVEYMEANS and PROC SURVEYREG) for calculating statistics based on specific sampling designs. Stata and SUDAAN are two other common statistical software packages that perform calculations for numerous statistics arising from the stratified, single-stage cluster sampling design.

NRD READMISSION RATES REPORTED ON THE HCUPNET WEBSITE

Readmission rates generated from the NRD are available on HCUPnet, a free online query system based on data from the HCUP (<u>http://hcupnet.ahrq.gov/</u>). This section describes how the readmission rates are calculated for HCUPnet to provide an example of how the NRD data elements might be used to define an index event, 7- and 30-day readmission, and readmission rates. Other types of readmission analyses are possible with the NRD; this is just one of many possible applications.

For the readmission rates on HCUPnet, an *index event* was defined as follows:

- Patient was discharged between January and November (1 <= DMONTH <= 11) for data years other than 2015.
 - For data year 2015, condition-specific readmission rates are based on index events from January to August with ICD-9-CM data.
- Patient was discharged alive (DIED = 0).
- Length of stay was non-missing (LOS >= 0).
- Discharge was for patient aged 1 year or older (AGE > 0).
- Patient may be a nonresident of the State (any value of RESIDENT).
- A patient is allowed to have multiple index events, regardless of how far apart.

For example, if a patient was discharged alive with a nonmissing length of stay on January 10, January 20, January 26, and March 30, all four discharges would qualify as index admissions.

For the readmission rates on HCUPnet, *readmissions* was defined as follows:

- The *first* discharge for a patient was within 7 or 30 days of an index event.
- Discharge occurred between January and December (1 <= DMONTH <= 12) for data years other than 2015.
 - For data year 2015, condition-specific readmission rates are based on readmissions from January to September with ICD-9-CM data.
- Discharge may be to the same or a different hospital (HOSP_NRD) and may result in a death.

¹⁵ Carlson BL, Johnson AE, Cohen SB. An evaluation of the use of personal computers for variance estimation with complex survey data. *J Off Statistics*. 1993;9(4):795-814.

On HCUPnet, the readmission rates were defined as the percentage of index admissions that had at least one readmission within 7 or 30 days.

- *Numerator* = total number of index events that had at least one subsequent hospital admission within 7 or 30 days.
- *Denominator* = total number of index events between January and November.
- *Rate* = numerator / denominator * 100.

Rates were not risk adjusted.

Consider an example of the 30-day, all-cause readmission rate for any diagnosis for a patient discharged alive on January 10, January 20, January 26, and March 30. Each admission is considered an index.

- January 10 is the first index admission.
- January 20 qualifies as a 30-day readmission for the January 10th index. It is also an index admission.
- January 26 qualifies as a 30-day readmission for the January 20th index. It is also an index admission.
- March 30 is an index admission, but it does not qualify as a readmission because it does not fall within 30 days of another index.

The 30-day readmission rate is 50 percent, because there are two 30-day readmissions for the four index admissions.

HCUPnet can be used to query 7- and 30-day readmission rates by various criteria. Queries can include the following criteria:

- Principal ICD-9-CM diagnosis using CCS (using the HCUP data element DXCCS1) through data year 2014
- All-listed ICD-9-CM external cause of injury CCS (using the HCUP data elements E_CCSn) through data year 2014
- All-listed ICD-9-CM major operating room procedures using CCS (using the HCUP data elements PRCCSn with the corresponding PCLASSn = 3 or 4) through data year 2014
- MDC (using the HCUP data element MDCnoPOA that does not consider the present on admission indicator for assignment)
- DRG (using the HCUP data element DRGnoPOA that does not consider the present on admission indicator for assignment).

HCUPnet reports readmission counts, rates, and costs stratified by characteristics of the index stay. Stratification characteristics include:

- Age group is based on the HCUP data element AGE.
- Sex is based on the HCUP data element FEMALE.
- Payer is assigned using the primary and secondary expected payer (HCUP data elements PAY1 and PAY2). If the primary or secondary expected payer indicates Medicare, then the payer category is assigned to Medicare. This categorization includes patients who are dually-eligible for Medicare and Medicaid under Medicare. If not

Medicare and the primary or secondary expected payer indicates Medicaid, then the payer category is Medicaid. If not Medicare or Medicaid and the primary or secondary expected payer indicates private insurance, then the payer category is Private. If not Medicare, Medicaid, or Private and the primary expected payer indicates self-pay or no charge, then the payer category is Self-pay/No charge. Stays for other types of payers are not reported on HCUPnet because this is a mixed payer group with small numbers. The expected secondary payer data element PAY2 is not available on the NRD.

- Income level is based on the HCUP data element ZIPINC_QRTL for the national quartile of the median household income for the patient's ZIP Code.
- Location is based on the HCUP data element PL_UR_CAT4 for the location of the
 patient's residence according to the Urban Influence Code (UIC) designation. Urban
 includes large and small metropolitan areas with all other areas categorized as rural.
 The data element PL_UR_CAT4 is not available on the NRD. The data element on the
 NRD for patient location is PL_NCHS, a six-category urban-rural classification scheme
 for U.S. counties developed by the National Center for Health Statistics (NCHS).

APPENDIX A: NRD INTRODUCTORY INFORMATION

| State | HCUP Data Source |
|----------------|---|
| Alaska | Alaska Department of Health and Social Services |
| Arkansas | Arkansas Department of Health |
| California | California Office of Statewide Health Planning and Developmen |
| Colorado | Colorado Hospital Association |
| Delaware | Delaware Division of Public Health |
| Florida | Florida Agency for Health Care Administration |
| Georgia | Georgia Hospital Association |
| Hawaii | Hawaii Laulima Data Alliance |
| Indiana | Indiana Hospital Association |
| Iowa | Iowa Hospital Association |
| Louisiana | Louisiana Department of Health |
| Maryland | Maryland Health Services Cost Review Commission |
| Massachusetts | Massachusetts Center for Health Information and Analysis |
| Mississippi | Mississippi Department of Health |
| Missouri | Missouri Hospital Industry Data Institute |
| Nebraska | Nebraska Hospital Association |
| Nevada | Nevada Department of Health and Human Services |
| New Mexico | New Mexico Department of Health |
| New York | New York State Department of Health |
| Oregon | Oregon Association of Hospitals and Health Systems |
| Pennsylvania | Pennsylvania Health Care Cost Containment Council |
| South Carolina | South Carolina Revenue and Fiscal Affairs Office |
| South Dakota | South Dakota Association of Healthcare Organizations |
| Tennessee | Tennessee Hospital Association |
| Utah | Utah Department of Health |
| Vermont | Vermont Association of Hospitals and Health Systems |
| Virginia | Virginia Health Information |
| Washington | Washington State Department of Health |
| Wisconsin | Wisconsin Department of Health Services |
| Wyoming | Wyoming Hospital Association |

Table A.1. HCUP Partners Participating in the 2019 NRD




Figure A.1. HCUP States Participating in the 2019 NRD

| Type of Discharge | Percentage of SID Discharges |
|--|---------------------------------|
| Included in the NRD | 83.6 |
| Excluded from the NRD | 16.4 |
| Hospital-level exclusions | |
| Noncommunity hospitals | 3.3 |
| Rehabilitation or LTAC hospitals | 0.2 |
| Discharge-level exclusions | |
| Discharges from all patients with an age of 0 (number of SID with excluded records) | 6.6 (14 of 30 SID) |
| Discharges with missing or unverified patient linkage numbers | 5.6 |
| Questionable patient linkage numbers: same patient linkage number on 20 or more discharges | 0.2 |
| Questionable patient linkage numbers: patient is hospitalized after discharged dead | 0.01 |
| Questionable patient linkage numbers: overlapping inpatient stays | 0.4 |
| Discharges from hospitals with more than 50 percent of their total discharges excluded for any of the above causes | 0.2 |

Table A.2. Percentage of SID Discharges in the NRD by Type of Discharge, 2019

| Year | States | Number of States for Discharges Aged 1 Year and Older | Number of States for Discharges Aged 0 | Number of Hospitals | Number of Discharges in the NRD, Unweighted | Number of Discharges in the NRD, Weighted |
|------|--|---|---|---------------------------|--|--|
| 2019 | AK, AR, CA, CO, DE, FL, GA, HI, IA, IN, LA, MA, MD, MO, MS, NE, NM, NV, NY, OR, PA, SC, SD, TN, UT, VA, VT, WA, WI, WY | 30 | 16 | 2,507 | 18,132,856 | 35,399,480 |
| 2018 | AK, AR, CA, DE, FL, GA, IA, IN, LA, MA, MD, MO, MS, NE, NM, NV, NY, OR, PA, SC, SD, TN, UT, VA, VT, WA, WI, WY | 28 | 14 | 2,430 | 17,686,511 | 35,460,557 |
| 2017 | AK, AR, CA, DE, FL, GA, IA, IN, LA, MA, MD, MO, MS, NE, NM, NV, NY, OR, PA, SC, SD, TN, UT, VA, VT, WA, WI, WY | 28 | 14 | 2,454 | 17,978,754 | 35,790,513 |
| 2016 | AK, AR, CA, FL, GA, HI, IA, LA, MA, MD, MO, MS, NE, NM, NV, NY, OR, PA, | 27 | 14 | 2,355 | 17,197,683 | 35,660,906 |

Table A.3. Summary of NRD States, Hospitals, and Inpatient Stays

| | SC, SD, TN, UT, VA, VT, WA, WI, WY | | | | | |
|------|---|----|----|-------|------------|------------|
| 2015 | AK, AR, CA, FL, GA, HI, IA, LA, MA, MD, MO, MS, NE, NM, NV, NY, OR, PA, SC, SD, TN, UT, VA, VT, WA, WI, WY | 27 | 14 | 2,367 | 17,198,125 | 35,673,252 |
| 2014 | AR, CA, FL, GA, HI, IA, LA, MA, MD, MO, NE, NM, NV, NY, SC, SD, TN, UT, VA, VT, WA, WI | 22 | 12 | 2,048 | 14,894,613 | 35,306,427 |
| 2013 | AR, CA, FL, GA, HI, IA, LA, MA, MO, NE, NM, NV, NY, SC, SD, TN, UT, VA, VT, WA, WI | 21 | 9 | 2,006 | 14,325,172 | 35,580,348 |
| 2012 | AK, AR, CA, FL, GA, HI, LA, MA, MO, NE, NM, NY, SC, TN, UT, VA, VT, WA | 18 | 6 | 1,715 | 13,459,216 | 36,465,049 |
| 2011 | AK, AR, CA, FL, GA, HI, LA, MA, MO, MS, NE, NM, NY, SC, TN, UT, VA, WA | 18 | 6 | 1,804 | 13,915,176 | 36,909,160 |
| 2010 | AK, AR, CA, FL, GA, HI, LA, MA, | 18 | 6 | 1,809 | 13,907,610 | 37,284,093 |

| MO, MS, NE, NM, | | | |
|---------------------------|--|--|--|
| NY, SC, TN, UT, VA, WA | | | |

Table A.4. NRD-Related Reports and Database Documentation Available on HCUP-US

| Description of the NRD Files | ICD-10-CM/PCS Data Included in the NRD Starting With 2015 | | | | |
|--|---|--|--|--|--|
| NRD Overview HCUP Partner in the NRD Introduction to the NRD, 2019 (<i>this document</i>) NRD Related Reports Restrictions on the Use HCUP Data Use Agreement Training Data Use Agreement for the HCUP Nationwide Databases Requirements for Publishing with HCUP data File Specifications and Load Programs | NRD Changes Beginning Data Year 2016 Caution: 2015 NRD Revised File Structure and New Data Elements Additional ICD-10-CM/PCS Resources HCUP Software Tools Tutorial Known Data Issues | | | | |
| NRD File Specifications – details data file names, number of records, record length, and record layout Nationwide SAS Load Programs Nationwide SPSS Load Programs Nationwide Stata Load Programs | Limitations on Using the NRD NRD, 2010-2012 NRD, 2010-2013 NRD, 2014 HCUP Tools: Labels and Formats | | | | |
| Data Elements in the NRD Availability of NRD Data Elements by Year – lists which data elements are available each year NRD Description of Data Elements – details uniform coding and State-specific idiosyncrasies Summary Statistics – lists means and frequencies on nearly all data elements Frequencies by Diagnosis and Procedure Codes, NRD 2016-2019 – includes frequency distributions for ICD-10-CM/PCS codes (individually and by the CCSR categories) | Clinical Classifications Software (CCS) Format Programs – to create value labels DRG Formats HCUP Formats HCUP Diagnoses and Procedure Groups Formats, including CCS Categories ICD-9-CM Formats ICD-10-CM/PCS Format Severity Formats | | | | |
| Additional Resources for Data Elements HCUP Quality Control Procedures – describes procedures used to assess data quality HCUP Coding Practices – describes how HCUP data elements are coded HCUP Hospital Identifiers – explains data elements that characterize individual hospitals | Obtaining HCUP Data Purchase HCUP Data from the HCUP Central Distributor | | | | |

| | Discharge Weights, 2019 | | |
|---------------------|-------------------------|--------------------------|---------|
| Patient Age and Sex | Minimum | Average Across Hospitals | Maximum |
| Males | | | |
| Age 0 | 1.08 | 4.58 | 23.25 |
| Age 1–17 | 1.10 | 3.54 | 23.89 |
| Age 18–44 | 1.03 | 1.97 | 9.29 |
| Age 45–64 | 1.04 | 1.93 | 9.04 |
| Age 65 and older | 1.04 | 1.94 | 8.78 |
| Females | | | |
| Age 0 | 1.07 | 4.57 | 24.81 |
| Age 1–17 | 1.09 | 2.92 | 15.06 |
| Age 18–44 | 1.02 | 1.93 | 8.66 |
| Age 45–64 | 1.03 | 1.90 | 8.51 |
| Age 65 and older | 1.05 | 1.93 | 8.54 |

Table A.5 Range of Discharge Weights by Patient Age and Sex, 2019

| Expected payer and age group ^a | Readmission rate | Number of readmissions ^b (thousands) |
|---|------------------|---|
| Medicare | | |
| Total ^c | 17.0 | 2,512 |
| 21–64 years | 21.3 | 601 |
| 65 years and above | 16.0 | 1,910 |
| Medicaid | | |
| Total | 13.8 | 849 |
| 1–20 years, non-maternal | 12.0 | 99 |
| 21–44 years, non-maternal | 17.6 | 277 |
| 45–64 years, non-maternal | 19.9 | 395 |
| Maternal | 4.4 | 79 |
| Private Insurance | | |
| Total | 8.6 | 594 |
| 1–20 years, non-maternal | 10.1 | 51 |
| 21–44 years, non-maternal | 10.0 | 139 |
| 45–64 years, non-maternal | 11.0 | 348 |
| Maternal | 3.1 | 58 |
| Self-pay/No charge ^d | | |
| Total | 11.8 | 144 |
| 1–20 years, non-maternal | 7.4 | 4 |
| 21–44 years, non-maternal | 11.9 | 66 |
| 45–64 years, non-maternal | 13.3 | 72 |
| Maternal | 4.0 | 3 |

 Table A.6 Rate and Number of 30-Day All-Cause Readmissions by Expected Payer and

 Patient Age Group, 2019

^a The expected payer is determined by the index admission, not the readmission. Information is not presented for three types of patients: (1) patients aged 0 years, (2) patients aged 1 year and above with other types of expected payer (e.g., local government programs, TRICARE, worker's compensation), and (3) patients aged 65 years and above with an expected payer of Medicaid, private insurance, or self-pay/no charge.

^b The number of readmissions are 12-month counts calculated by multiplying the average readmission rate by the 12month index admission counts. The sum of the age groups for each payer category may not equal the total as the result of rounding.

^c Medicare patients under age 21 years are included in the Medicare total for all ages but are not reported separately because of the small number of cases. There is no maternal breakout for Medicare because of the small number of maternal stays in this group.

^d Self-pay/no charge patients have an expected primary payer of self-pay, no charge, charity, or no expected payment

 Table A.7 Rate and Number of 30-Day All-Cause Readmissions by the 20 Most Common

 Principal Diagnoses at Index Admission, Ranked by Number of Readmissions, 2019

| Rank | Principal Diagnosis at Index Admission | Number of 30- Day All-Cause Readmissions | 30-Day All-Cause Readmission Rate ^a |
|------|---|--|--|
| 1 | INF002: Septicemia | 325,758 | 17.5 |
| 2 | CIR019: Heart failure | 238,343 | 22.7 |
| 3 | END003: Diabetes mellitus with complication | 126,520 | 19.8 |
| 4 | RSP008: Chronic obstructive pulmonary disease and bronchiectasis | 100,203 | 19.9 |
| 5 | RSP002: Pneumonia (except that caused by tuberculosis) | 97,622 | 15.3 |
| 6 | GEN002: Acute and unspecified renal failure | 96,317 | 19.4 |
| 7 | RSP012: Respiratory failure; insufficiency; arrest | 84,767 | 20.7 |
| 8 | CIR017: Cardiac dysrhythmias | 84,105 | 14.3 |
| 9 | MBD001: Schizophrenia spectrum and other psychotic disorders | 83,807 | 21.6 |
| 10 | FAC006: Encounter for antineoplastic therapies | 83,416 | 68.1 |
| 11 | CIR009: Acute myocardial infarction | 74,480 | 13.5 |
| 12 | GEN004: Urinary tract infections | 72,428 | 15.4 |
| 13 | END011: Fluid and electrolyte disorders | 63,336 | 20.0 |
| 14 | MBD002: Depressive disorders | 62,175 | 11.9 |
| 15 | INJ037: Complication of other surgical or medical care, injury, initial encounter | 61,150 | 19.5 |
| 16 | MBD017: Alcohol-related disorders | 60,637 | 19.3 |
| 17 | SKN001: Skin and subcutaneous tissue infections | 55,928 | 12.2 |
| 18 | CIR020: Cerebral infarction | 54,473 | 11.3 |
| 19 | DIG021: Gastrointestinal hemorrhage | 52,327 | 18.2 |
| 20 | GEN003: Chronic kidney disease | 48,366 | 28.2 |

Notes: Principal diagnosis category is based on the <u>Clinical Classifications Software Refined (CCSR)</u> default categorization scheme for the principal diagnosis in inpatient data, v2021.2. The weighted number of 30-day all-cause readmissions is presented for the top 20 principal diagnoses at the index stay. Diagnoses are sorted by the weighted number of readmissions. The definitions of an index admission and readmission are consistent with <u>HCUPnet</u>.

^a Rate per 100 Index Stays

Table A.8 Rate and Number of 30-Day All-Cause Readmissions by Major Diagnostic Category(MDC) at Index Admission, Ranked by Number of Readmissions, 2019

| Rank | Major Diagnostic Category (MDC) | Number of All- Cause, 30-Day Readmissions | All-Cause, 30- Day Readmission Rate ^a |
|------|---|---|--|
| 1 | 5: Diseases & Disorders of The Circulatory System | 709,641 | 17.3 |
| 2 | 4: Diseases & Disorders of The Respiratory System | 442,788 | 16.6 |
| 3 | 6: Diseases & Disorders of The Digestive System | 389,488 | 15.5 |
| 4 | 18: Infectious & Parasitic Diseases, Systemic or Unspecified Sites | 361,427 | 17.3 |
| 5 | 11: Diseases & Disorders of The Kidney & Urinary Tract | 267,254 | 17.8 |
| 6 | 8: Diseases & Disorders of The Musculoskeletal System & Conn Tissue | 250,727 | 8.2 |
| 7 | 1: Diseases & Disorders of The Nervous System | 247,507 | 13.0 |
| 8 | 19: Mental Diseases & Disorders | 223,035 | 15.4 |
| 9 | 10: Endocrine, Nutritional & Metabolic Diseases & Disorders | 188,212 | 16.1 |
| 10 | 7: Diseases & Disorders of The Hepatobiliary System & Pancreas | 176,179 | 18.9 |
| 11 | 14: Pregnancy, Childbirth & The Puerperium | 135,093 | 3.8 |
| 12 | 17: Myeloproliferative Diseases & Disorders, Poorly Differentiated Neoplasm | 123,265 | 49.6 |
| 13 | 16: Diseases & Disorders of Blood, Blood Forming Organs, Immunological Disorders | 100,487 | 24.5 |
| 14 | 9: Diseases & Disorders of The Skin, Subcutaneous Tissue & Breast | 84,188 | 12.6 |
| 15 | 20: Alcohol/Drug Use & Alcohol/Drug Induced Organic Mental Disorders | 79,289 | 17.7 |
| 16 | 21: Injuries, Poisonings & Toxic Effects of Drugs | 61,463 | 13.9 |
| 17 | 23: Factors Influencing Health Status & Other Contacts with Health Services | 28,804 | 18.1 |
| 18 | 3: Diseases & Disorders of The Ear, Nose, Mouth & Throat | 23,426 | 9.4 |
| 19 | 13: Diseases & Disorders of The Female Reproductive System | 20,311 | 8.3 |
| 20 | 12: Diseases & Disorders of The Male Reproductive System | 10,348 | 9.8 |
| 21 | 24: Multiple Significant Trauma | 7,907 | 9.4 |
| 22 | 25: Human Immunodeficiency Virus Infections | 7,832 | 24.1 |
| 23 | 2: Diseases & Disorders of The Eye | 3,444 | 8.9 |
| 24 | 22: Burns | 2,436 | 9.1 |

Notes: MDC 0, Principal DX Cannot be Assigned to MDC, is not included. The definitions of an index admission and readmission are consistent with <u>HCUPnet</u>.

^a Rate per 100 Index Stays

APPENDIX B: HANDLING OF TRANSFERS AND SAME-DAY STAYS

Hospital administrative databases like the NRD and SID are "discharge-level" files, meaning that each record represents one discharge abstract from an inpatient stay. If a patient visits the hospital multiple times in a given year, the SID includes separate records for each inpatient stay. In addition, if a patient is transferred between hospitals within the State, the SID contain two discharge records, one record from the first hospitalization and a second record from the latter hospitalization.¹⁶

Defining Transfers and Same-Day Events

Readmission analyses do not usually allow the hospitalization at the receiving hospital to be counted as a readmission. To eliminate this possibility, the pairs of records representing a transfer were collapsed into a single "combined" record in the NRD and removed the original separate discharge records from the NRD. *Transfer* records were defined as having all of the following characteristics:

- Discharge date of the first inpatient stay equaled the admission date of a subsequent inpatient stay.¹⁷
- The first record had a discharge disposition of transfer to an acute care hospital.
- The second record was from a different hospital and had an admission source indicating a transfer.

A discharge was defined as a *same-day* stay if the discharge date for one inpatient stay was the same as the admission date of a second stay for the same patient (same as transfers), but there was no indication of a transfer by the discharge disposition or admission source. Same-day stays may or may not have involved different hospitals. Same-day stays may indicate that a patient was discharged too soon and then needed to be returned to the hospital on the same day. However, it was also possible that these were transfer records with an incorrect or missing discharge disposition and admission source.

Records that were part of transfers or same-day stays were collapsed into a single combined record. These combined records account for about three percent of records in the NRD and are identified by the data element SAMEDAYEVENT. The value of SAMEDAYEVENT is defined as follows:

- Transfer involving two discharges from <u>different</u> hospitals (value 1): Discharge date of one admission equaled the admission date on another record for the same patient. There was a discharge disposition of transfer to an acute care hospital on the first record and an admission source of transfer (in) on the subsequent record; two different hospitals were involved.
- Same-day stay involving two discharges at <u>different</u> hospitals (value 2): The discharge date on the first inpatient stay equaled the admission date on a second inpatient stay for the same patient; two different hospitals were involved, but the coding of either the discharge disposition or admission source did not indicate a transfer.

 ¹⁶ If the patient is transferred to an out-of-State hospital, the subsequent discharge would not be included in the NRD because the HCUP patient linkage numbers only can follow a patient within a State.
 ¹⁷ Although the text refers to using the discharge and admission date, in reality, the NRD data elements

NRD_DaysToEvent and LOS were used to identify the sequential order of inpatient records and whether they stopped and started on the same day.

- Same-day stays involving two discharges at the <u>same</u> hospital (value 3): The discharge date on the first inpatient stay equaled the admission date on a second inpatient stay for the same patient; the hospital was the same on both records.
- Same-day stay involving three or more discharges, same or different hospitals (value 4): Multiple records indicating a combination of transfers and other same-day events seem odd, but they do rarely occur in administrative data.
- Not a transfer or other same-day stay (value 0). About 97 percent of the records in the NRD did not involve transfers or same-day stays.

Please note that if a patient had a discharge disposition of transfer and an admission source that also indicated a transfer, but the discharge date of the first stay did not equal the admission date of the second stay (e.g., the patient was admitted the next day because the transfer occurred at night), the two records are not considered a transfer in the NRD.

Creating a Combined Transfer Record

Combining information across transfer and same-day stay records required specific rules for how to handle different types of information on the pairs of records. The pairs of records were first ordered by the earliest occurrence in the year. The different scenarios described below detail how different types of information were combined:

- Use first. For information that pertains to day of admission (e.g., admission month, admission source, NRD_DaysToEvent), the combined transfer record used the information from the first record in the pair.
- Use last record. For information that pertains to the end of the time in the hospital (e.g., discharge disposition, expected payer, , the combined transfer record used the information from the latter record in the pair.
- Summarize. For information that needed to reflect both stays, the combined transfer record summarized the information. For example, length of stay and total charge were summed across the pair of records.
- Diagnoses and related data elements. Each record included arrays of ICD-9-CM or ICD-10-CM diagnoses and related data elements. On the assumption that the diagnoses on the latter record would reflect the final determination of diagnoses after complete treatment, the diagnoses reported on the latter record were retained at the beginning of the diagnosis array (including the principal diagnosis). The diagnoses from the first record was added to the end of the diagnosis array. Diagnosis codes in the NRD are limited to a maximum of 25 (prior to 2014), 30 (starting in 2014), and 40 (starting in 2017) with no more than the maximum number of diagnoses retained on a combined transfer record. Less than one percent of diagnoses are truncated. The same scheme was used for all diagnosis-related data elements also contained in arrays, including diagnosis-related group (DRG) and major diagnostic category (MDC).
 - Exception for handling transfers related to rehabilitation or after care (when the transfer records used the same diagnosis/procedure coding system [ICD-9-CM or ICD-10-CM/PCS]): In some cases, there may be an indication that a patient was transferred for rehabilitation or aftercare. Using the principal diagnosis from the second part of the stay on the combined record would mask the real reason for the admission. Therefore, the assignment of diagnoses and related data elements (including DRG and MDC) was modified to give precedence to the information on the first inpatient stay with the information from

the latter stay added to the end of the associated arrays. The criteria to identify rehabilitation- or aftercare-related records has changed over time:

Beginning data year 2018

An indication of rehabilitation or aftercare is identified based on either of the following:

- Any record but the last record had a discharge disposition of 62 (Discharged/transferred to an Inpatient Rehabilitation Facility [IRF] including Rehabilitation Distinct part units of a hospital) or 90 Discharged/Transferred to an Inpatient Rehabilitation Facility (IRF) including Rehabilitation Distinct Part Units of a Hospital with a Planned Acute Care Hospital Inpatient Readmission (Effective 10/1/13).
- Any record but the first record had default CCSR FAC010, Other aftercare encounter, for the principal diagnosis.

Through data year 2017

An indication of rehabilitation or aftercare was identified if the latter record had any one of the following Clinical Classifications Software (CCS) categories:

- CCS 254: Rehabilitation care; fitting of prostheses; and adjustment of devices
- CCS 255: Administrative/social admission
- CCS 256: Medical examination/evaluation
- CCS 257: Other aftercare
- CCS 258: Other screening for suspected conditions (not mental disorders or infectious disease)
- CCS 259: Residual codes; unclassified.

These types of transfer records are flagged by the data element REHABTRANSFER (value 1).

- Exception to this rule when the transfer records included different diagnosis/procedure coding system: If the first record included ICD-9-CM codes and the latter record was discharged after October 1, 2015 with ICD-10-CM/PCS codes, then only the ICD-10-CM/PCS diagnosis and procedure codes were retained on the combined transfer record.
- Procedures and related data elements. Each record included arrays of ICD-9-CM or ICD-10-PCS procedure codes and related data elements. Although procedure codes were not necessarily reported in the order in which they were performed, it was decided to retain the procedures reported on the latter record at the beginning of the procedure array to mirror the approach for diagnoses. The procedures from the first record were added to the end of the procedure array. The NRD is limited to a maximum of 15 procedure codes (prior to 2017) and 25 procedure codes (starting in 2017), so no more than the maximum number of procedures were retained on a combined transfer record. The same scheme was used for all procedure-related data elements that were also contained in arrays. The day of procedure on the latter record was adjusted for the length of stay on the first record, so the days reflected the time from the beginning of the first stay.

- *Hospital identifiers and characteristics*. The hospital identifiers and characteristics from the latter stay were retained on the combined transfer record. This assumed that the latter hospital was primarily responsible for the care.
- *Patient characteristics*. The patient characteristics (e.g., age, sex, income quartile) from the first stay were retained on the combined transfer record.

APPENDIX C: STATE-SPECIFIC RESTRICTIONS

The table below enumerates the types of restrictions applied to the Nationwide Readmissions Database. Restrictions include the following types:

- Confidentiality of records
 - Restricted release of age in years
- Missing discharges for specific populations of patients.

Table C.1. State-Specific Restrictions

Confidentiality of Records

Restricted release of patient's age in years to ensure patient confidentiality:

 Age (AGE) values greater than 90 are aggregated into a single category of 90 years or older in the NRD.

Missing Discharges for Specific Populations of Patients

The following data sources may be missing discharge records for specific populations of patients:

- At least one Partner prohibits the release of discharge records for patients with HIV diagnoses.
- At least one Partner prohibits the release of inpatient records that were discharged from Alternative Level of Care units of the hospital (e.g., skilled nursing or swing bed units).
- At least one Partner prohibits the release of abortion discharges.
- At least one Partner restricts the release of ICD-10-CM diagnosis codes for medical misadventures and adverse effects.

APPENDIX D: NRD DATA ELEMENTS AND CODES

| Category | Data Element Name | Description |
|-------------------------|-------------------|---|
| | AWEEKEND | Admission on weekend: 0) admission on Monday–Friday, (1) admission on Saturday– Sunday |
| | DIED | Indicates in-hospital death: 0) did not die during hospitalization, (1) died during hospitalization |
| Admission/ | DISPUNIFORM | Disposition of patient, uniform coding: (1) routine, (2) transfer to short term hospital, (5) other transfers, including skilled nursing facility, intermediate care, and another type of facility, (6) home health care, (7) against medical advice, (20) died in hospital, (99) discharged alive, destination unknown |
| Discharge | DQTR | Coded: (1) Jan–Mar, (2) Apr–Jun, (3) Jul–Sep, (4) Oct–Dec |
| | ELECTIVE | Indicates elective admission: (1) elective, (0) nonelective admission |
| | HCUP_ED | Indicator that discharge record includes evidence of emergency department (ED) services: (0) record does not meet any HCUP ED criteria, (1) ED revenue code was on SID record, (2) ED charge reported on SID record, (3) ED CPT procedure code on SID record, (4) other indication of ED services |
| | YEAR | Discharge year |
| | DRG | DRG in use on discharge date |
| | DRG_NoPOA | DRG assignment made without the use of the present on admission flags for the diagnoses |
| | DRGVER | Grouper version in use on discharge date |
| | I10_BIRTH | ICD-10-CM birth indicator |
| Clinical Information | I10_DELIVERY | ICD-10-CM/PCS delivery indicator |
| | I10_DX1-I10_DX40 | ICD-10-CM diagnoses, principal and secondary |
| | I10_INJURY | Injury ICD-10-CM diagnosis reported on record |
| | I10_MULTINJURY | Multiple ICD-10-CM injuries reported on record |
| | I10_NDX | Number of ICD-10-CM diagnoses coded on the record |

Table D.1. Data Elements in the 2019 NRD Core File

| Category | Data Element Name | Description |
|--------------------|-------------------|---|
| | I10_NPR | Number of procedures coded |
| | I10_PR1-I10_PR25 | ICD-10-PCS procedures, principal and secondary |
| | I10_SERVICELINE | ICD-10-CM/PCS hospital service line indicator |
| | MDC | MDC in use on discharge date |
| | MDC_NoPOA | MDC assignment made without the use of the present on admission flags for the diagnoses |
| | PCLASS_ORPROC | ICD-10-PCS major operating room procedure |
| | PRDAY1-PRDAY25 | The day on which the procedure is performed. A value of 0 indicates the day of admission. |
| | HOSP_NRD | NRD hospital identifier specific to the NRD and is not linkable to any other HCUP or external databases. HOSP_NRD can be used to add data elements from the Hospital file to records on the discharge-level files. The values of HOSP_NRD differ from year to year. An individual hospital cannot be tracked across data years. |
| NRD Identifiers | KEY_NRD | Unique record identifier for the discharge in the NRD and not linkable to any other HCUP or external databases. KEY_NRD can be used to add data elements from the Severity and Diagnosis/Procedure Groups files to the records on the Core file within the same data year. The values of KEY_NRD are different in each data year 2010–2012 and 2015–2019 but are nonunique between 2013 and 2014. |
| | | Please note that KEY_NRD is a record identifier and not a patient linkage number. NRD_VISITLINK is the patient linkage number specific to the NRD. |
| Patient | AGE | Age in years coded 0-90 years; any age greater than 90 was set to 90. Missing age was imputed using other records with the same patient linkage number. In the 2019 NRD, about 189 discharges (0.00 percent) had the age imputed. |
| Demographics | FEMALE | Indicates sex: (0) male, (1) female. Missing sex was imputed using other records with the same patient linkage number. In the 2019 NRD, about 122 discharges (0.00 percent) had the sex imputed. |

| Category | Data Element Name | Description |
|-------------------------|-------------------|---|
| | PAY1 | Expected primary payer, uniform: (1) Medicare, (2) Medicaid, (3) private insurance, (4) self-pay, (5) no charge, (6) other |
| | PL_NCHS | Patient location: National Center for Health Statistics (NCHS) urban-rural classification scheme for U.S. counties: (1) "Central" counties of metro areas of >=1 million population, (2) "Fringe" counties of metro areas of >=1 million population, (3) Counties in metro areas of 250,000–999,999 population, (4) Counties in metro areas of 50,000–249,999 population, (5) Micropolitan counties, (6) Not metropolitan or micropolitan counties |
| | ZIPINC_QRTL | Median household income quartiles for patient's ZIP Code: (1) quartile 1 [lowest income], (2) quartile 2, (3) quartile 3, (4) quartile 4 [highest income]. For 2019, the median income quartiles are defined as: (1) \$1-\$45,999; (2) \$46,000-\$58,999; (3) \$59,000-\$78,999; and (4) \$79,000 or more. |
| | DMONTH | Discharge month coded from (1) January to (12) December |
| | NRD_DAYSTOEVENT | Count of days from randomly selected "start date" to admission date coded differently for each value of NRD_VisitLink |
| Readmission Specific | NRD_VISITLINK | Patient linkage number specific to the NRD and not linkable to any other HCUP or external databases. The values of NRD_VISITLINK differ from year to year. An individual person cannot be tracked across data years. |
| | REHABTRANSFER | A combined record involving transfer to rehabilitation, evaluation, or other aftercare: (1) yes, (0) no |
| | RESIDENT | Identifies patient as a resident of the State in which he or she received hospital care: (1) resident, (0) nonresident |

| Category | Data Element Name | Description |
|-----------|-------------------|--|
| | SAMEDAYEVENT | Identifies records that were combined from transfer or same-day stay pairs of records: (0) not a combined transfer or other same-day stay record, (1) combined transfer involving two discharges from different hospitals, (2) combined same-day stay involving two discharges at different hospitals, (3) combined same-day stay involving two discharges at the same hospital, (4) combined same-day stay involving three or more discharges at same or different hospitals |
| Resource | LOS | Length of stay, edited |
| Use | ТОТСНС | Total charges, edited |
| | DISCWT | NRD discharge weight to be used for calculating national estimates |
| Weighting | NRD_STRATUM | NRD stratum for post-stratification based on geographic region, urban/rural location, teaching status, size of hospital based on number of beds, and control/ownership. For the confidentiality of hospitals and States, the NRD_STRATUM was randomly assigned. The values of NRD_STRATUM differ from year to year. An individual stratum cannot be tracked across data years. |

Table D.2. Data Elements in the 2019 NRD Severity Measures File

For prior years, refer to the <u>NRD Description of Data Elements</u> page on the HCUP-US website or to previous versions of the NRD Introduction.

| Category | Data Element Name | Description |
|--------------------|-----------------------|--|
| | APRDRG | 3M All Patient Refined DRG |
| 3M | APRDRG_Risk_Mortality | 3M All Patient Refined DRG: Risk of Mortality Subclass: (0) No class specified, (1) Minor likelihood of dying, (2) Moderate likelihood of dying, (3) Major likelihood of dying, (4) Extreme likelihood of dying |
| APR-DRG | APRDRG_Severity | 3M All Patient Refined DRG: Severity of Illness Subclass: (0) No class specified, (1) Minor loss of function (includes cases with no comorbidity or complications), (2) Moderate loss of function, (3) Major loss of function, (4) Extreme loss of function |
| NRD Identifiers | HOSP_NRD | NRD hospital identifier specific to the NRD and is not linkable to any other HCUP or external databases. HOSP_NRD can be used to add data elements from the Hospital file to records on the discharge-level files. The values of HOSP_NRD differ from year to year. An individual hospital cannot be tracked across data years. |
| | KEY_NRD | Unique record identifier for the discharge in the NRD and not linkable to any other HCUP or external databases. KEY_NRD can be used to add data elements from the Severity and Diagnosis/Procedure Groups files to the records on the Core file within the same data year. The values of KEY_NRD are different in each data year 2010–2012 and 2015–2019 but are not unique between 2013 and 2014. |
| | | Please note that KEY_NRD is a record identifier and not a patient linkage number. NRD_VISITLINK is the patient linkage number specific to the NRD. |

Table D.3. Data Elements in the 2019 NRD Diagnosis and Procedure Groups Files

For data years 2016-2017, the Diagnosis and Procedure Groups File is not available on the NRD.

For data 2014 and prior years, refer to the <u>NRD Description of Data Elements</u> page on the HCUP-US website or to previous versions of the NRD Introduction.

| Category | Data Element Name | Description |
|---------------------------------------|----------------------------|--|
| Elixhauser Comorbidity Software | CMR_aaa ¹ | Comorbidity measures (aaa) identified by the AHRQ Elixhauser Comorbidity Software Refined for ICD-10-CM diagnosis codes |
| Refined for ICD-10-CM | CMR_VERSION | Version of the Elixhauser Comorbidity Measure Refined for ICD-10-CM |
| | DXCCSR_aaannn ² | Indication that at least one ICD-10-CM diagnosis on the record is included in CCSR aaannn |
| Clinical Classifications | DXCCSR_DEFAULT_DX1 | Default CCSR for principal/first-listed ICD-10- CM diagnosis |
| Software Refined | DXCCSR_VERSION | Version of CCSR for ICD-10-CM diagnoses |
| (CCSR) Category | PRCCSR_aaannn ³ | Indication that at least one ICD-10-PCS procedure code on the record is included in CCSR aaannn |
| | PRCCSR_VERSION | Version of the CCSR for ICD-10-PCS procedures |
| Procedure | PCLASSn ^₄ | Procedure Classes Refined for ICD-10-PCS procedures |
| Classes Refined for | PCLASS_ORPROC | ICD-10-PCS major operating room procedure indicator |
| ICD-10-PCS | PCLASS_VERSION | Version of the Procedure Classes Refined for ICD-10-PCS procedures |
| NRD Identifiers | HOSP_NRD | NRD hospital identifier specific to the NRD and is not linkable to any other HCUP or external databases. HOSP_NRD can be used to add data elements from the Hospital file to records on the discharge-level files. The values of HOSP_NRD differ from year to year. An individual hospital cannot be tracked across data years. |

¹ Where aaa denotes the specific comorbidity measure.

² Where aaa denotes the body system and nnn denotes the CCSR number within the body system.

³ PCLASSn was also available on the NRD through quarter 3 of data year 2015 and was specific to the coding of ICD-9-CM procedures.

⁴ Where aaa denotes the clinical domain and nnn denotes the CCSR number within the clinical domain.

| Category | Data Element Name | Description |
|----------|-------------------|--|
| | KEY_NRD | Unique record identifier for the discharge in the NRD and not linkable to any other HCUP or external databases. KEY_NRD can be used to add data elements from the Severity and Diagnosis/Procedure Groups files to the records on the Core file within the same data year. The values of KEY_NRD are different in each data year 2010–2012 and 2015–2019 but are nonunique between 2013 and 2014. Please note that KEY_NRD is a record identifier and not a patient linkage number. NRD_VISITLINK is the patient linkage number specific to the NRD. |

| Category | Data Element Name | Description |
|-------------------------|-------------------|---|
| Admission/ Discharge | YEAR | Discharge year |
| | H_CONTROL | Control/ownership of hospital: (1) government, nonfederal [public], (2) private, not-for-profit [voluntary], (3) private, investor-owned [proprietary] |
| | HOSP_BEDSIZE | Size of hospital based on the number of beds: (1) small, (2) medium, (3) large. The categories are defined using region of the U.S., the urban-rural designation of the hospital, in addition to the teaching status. |
| | HOSP_UR_TEACH | Teaching status of hospital: (0) metropolitan non- teaching, (1) metropolitan teaching, (2) non- metropolitan |
| Hospital Information | HOSP_URCAT4 | Hospital urban-rural location: (1) large metropolitan areas with at least 1 million residents, (2) small metropolitan areas with less than 1 million residents, (3) micropolitan areas, (4) not metropolitan or micropolitan, (8) metropolitan, collapsed category of large and small metropolitan, (9) non-metropolitan, collapsed category of micropolitan and rural |
| | NRD_STRATUM | NRD stratum for post-stratification based on geographic region, urban/rural location, teaching status, bed size, and control. Region is not identified. The values of NRD_STRATUM differ from year to year. An individual stratum cannot be tracked across data years. |
| NRD Identifiers | HOSP_NRD | NRD hospital identifier specific to the NRD and is not linkable to any other HCUP or external databases. The values of HOSP_NRD differ from year to year. An individual hospital cannot be tracked across data years. |
| | N_DISC_U | Number of discharges in the target universe in the stratum |
| | N_HOSP_U | Number of hospitals in the target universe in the stratum |
| Weighting | S_DISC_U | Number of NRD discharges in the stratum |
| | S_HOSP_U | Number of NRD hospitals in the stratum |
| | TOTAL_DISC | Total number of discharges for this hospital in the NRD |

 Table D.4. Data Elements in the 2019 NRD Hospital File

APPENDIX E: EVALUATION OF THE DIFFERENCE IN READMISSION RATES CAUSED BY HCUP PATIENT LINKAGE NUMBERS BEING SPECIFIC TO A STATE

The HCUP revisit variables (NRD_VisitLink and NRD_DaysToEvent) that can be used to track patients across hospitalizations were created from patient linkage numbers provided by the HCUP Partners. These identifiers can only track patients across hospitals in a single State. Consider the following two illustrative examples:

- 1. If a person is hospitalized in February in New York and then hospitalized in April to the same hospital or to a different hospital in New York, the HCUP revisit variables for the New York SID will be able to track the patient across the two visits.
- 2. If a person is hospitalized in February in New York and then hospitalized in April to a hospital in Florida, the HCUP revisit variables for the New York SID will not track the patient between New York and Florida.

The HCUP revisit variables are specific to tracking patients hospitalized in a State, regardless of whether the patient is a resident of the State. This limitation in the HCUP revisit variables causes the readmission rates to be artificially low. Consider the following example:

- Patient A has two index events in California and was readmitted one time in California within 30 days.
- Patient B has three index events in California but was in Florida when readmitted within 30 days.

The true readmission rate across these two patients is 0.40 (0.40 = 2 readmissions / 5 index events). Using only California data, the readmission rate is 0.20 (0.20 = 1 readmission / 5 index events). Using California and Florida data combined, the readmission rate is 0.17 (0.17 = 1 readmission / 6 index events); there are 6 index events because the readmission for patient B in Florida gets counted as an index event because it cannot be tied to the hospitalization in California).

The 2011 Medicare Standard Analytic File (SAF) was used to examine the impact on readmission rates caused by having State-specific patient linkage numbers. The SAF includes patient linkage numbers that follow Medicare Fee-For-Service (FFS) patients across States; therefore, they do not have the same limitations as the HCUP data. Condition-specific readmission rates were calculated in two ways. The index event for both readmission rates was allowed to include resident and nonresident discharges, similar to the NRD. The index events were grouped by the AHRQ Clinical Classifications Software (CCS) category for the principal diagnosis. For the first set of readmission rates, the 30-day readmission for any cause was required to occur in the same State as the index event, similar to the NRD. For the second set of readmission rates, was limited to CCS categories with at least 100 index events. The comparison of readmission rates was limited to CCS categories with at least 100 index events. There were 247 CCS categories and the number of index events ranged from 101 to 468,709.

The analysis of readmission rates using the 2011 Medicare FFS data demonstrated that condition-specific readmission rates were higher if a patient could be tracked across all States, but that the percentage increase was less than 5 percent for most of the CCS categories. This analysis was limited in that it focused on the Medicare FFS population. The Medicare population accounted for 40 percent of all inpatient discharges in 2011,¹ and previous research indicates that this population has higher readmission rates than discharges for other payers.²

¹ HCUPnet query on the expected primary payer for the 2011 Nationwide Inpatient Sample. Accessed December 9, 2014. ² Wier LM, Barrett ML, Steiner C, Jiang HJ. All-Cause Readmissions by Payer and Age, 2008. HCUP Statistical Brief #115. June 2011. Rockville, MD: Agency for Healthcare Research and Quality.

Conditions often associated with younger adults, such as pregnancy, were included in the Medicare estimates because 20 percent of Medicare discharges are under the age of 65.³ Medicare patients under the age of 65 include people who are disabled or who have been diagnosed with end-stage renal disease or amyotrophic lateral sclerosis (ALS). Given the volume and severity of illness for Medicare patients, the estimates for the increase in the condition-specific readmission rates using the Medicare data provided a reasonable upper bound on the impact. The following three tables provide more detail on specific changes in readmission rates.

Table E.1 lists the percentage increase between the two types of readmission rates for the 20 conditions with the largest number of index events. These 20 conditions represent 54 percent of the index events in the analysis. The increase in the readmission rates when readmissions were captured in other States ranged from 1.9 percent for urinary tract infections to 3.7 percent for chest pain.

| Table E.1. Twenty Conditions with the Largest Number of Index Events, Restricting |
|---|
| Readmissions to Within State and Across All States |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|---------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 108: Congestive heart failure; nonhypertensive | 468,709 | 25.05 | 25.70 | 2.6 |
| 122: Pneumonia (except that caused by tuberculosis or sexually transmitted disease) | 443,388 | 18.25 | 18.69 | 2.4 |
| 2: Septicemia (except in labor) | 405,898 | 21.42 | 21.87 | 2.1 |
| 203: Osteoarthritis | 363,295 | 5.57 | 5.70 | 2.2 |
| 127: Chronic obstructive pulmonary disease and bronchiectasis | 356,562 | 21.99 | 22.40 | 1.9 |
| 106: Cardiac dysrhythmias | 345,225 | 16.73 | 17.27 | 3.3 |
| 159: Urinary tract infections | 277,674 | 18.31 | 18.66 | 1.9 |

³ HCUPnet query on the expected primary payer and age for the 2011 Nationwide Inpatient Sample. Accessed December 9, 2014.

| | | | mission Rate dex Events | Percentage Increase in | |
|---|---------------------------------|---|---|--|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States | |
| 237: Complication of device; implant or graft | 274,669 | 21.23 | 21.83 | 2.8 | |
| 109: Acute cerebrovascular disease | 232,510 | 14.73 | 15.23 | 3.4 | |
| 157: Acute and unspecified renal failure | 228,134 | 22.04 | 22.60 | 2.5 | |
| 101: Coronary atherosclerosis and other heart disease | 220,481 | 15.44 | 15.99 | 3.5 | |
| 100: Acute myocardial infarction | 205,949 | 19.66 | 20.41 | 3.8 | |
| 55: Fluid and electrolyte disorders | 192,425 | 20.33 | 20.81 | 2.4 | |
| 205: Spondylosis; intervertebral disc disorders; other back problems | 178,614 | 10.62 | 10.93 | 2.9 | |
| 197: Skin and subcutaneous tissue infections | 178,097 | 16.09 | 16.45 | 2.2 | |
| 226: Fracture of neck of femur (hip) | 172,566 | 13.89 | 14.19 | 2.2 | |
| 50: Diabetes mellitus with complications | 159,421 | 23.81 | 24.31 | 2.1 | |
| 238: Complications of surgical procedures or medical care | 155,584 | 20.50 | 21.08 | 2.9 | |
| 153: Gastrointestinal hemorrhage | 155,410 | 18.75 | 19.27 | 2.8 | |
| 102: Nonspecific chest pain | 148,512 | 13.50 | 14.00 | 3.7 | |

Table E.2 lists the 10 conditions with the largest percentage differences between the two types of readmission rates. The readmission rates using discharges across all States were expected to be higher than the readmissions restricted to the same State as the index event. Only three CCS categories had an increase of more than 10 percent in the two readmission rates, and these CCS categories had a very small number of index events. The other seven CCS categories had a percentage increase between 6.6 percent and 7.9 percent and also had relatively small numbers of index events.

Table E.2. Ten Conditions with the Largest Difference in Readmission Rates, Restricting Readmissions to Within State and Across All States

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|---------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 188: Fetopelvic disproportion; obstruction | 153 | 4.58 | 5.23 | 14.3 |
| 20: Cancer; other respiratory and intrathoracic | 952 | 17.65 | 19.85 | 12.5 |
| 655: Mental disorders usually diagnosed in infancy, childhood, or adolescence | 115 | 15.65 | 17.39 | 11.1 |
| 227: Spinal cord injury | 2,420 | 17.69 | 19.09 | 7.9 |
| 185: Prolonged pregnancy | 665 | 2.11 | 2.26 | 7.2 |
| 77: Encephalitis (except that caused by tuberculosis or sexually transmitted disease) | 1,945 | 17.22 | 18.46 | 7.2 |
| 240: Burns | 4,110 | 16.40 | 17.54 | 7.0 |
| 213: Cardiac and circulatory congenital anomalies | 2,502 | 13.75 | 14.71 | 7.0 |
| 670: Miscellaneous disorders | 3,592 | 13.34 | 14.23 | 6.7 |
| 96: Heart valve disorders | 47,909 | 21.73 | 23.16 | 6.6 |

Table E.3 has the complete listing of readmissions rates for the 247 CCS categories. The majority of CCS categories (221 of the 247, 89.5 percent) had a percentage increase between the two types of readmission rates of less than 5 percent.

Table E.3. Readmission Rates Restricting Readmissions to Within State and Across All States, Medicare Standard Analytic File, 2011

| | | | mission Rate lex Events | Percentage Increase in |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 1: Tuberculosis | 955 | 21.99 | 22.72 | 3.3 |
| 2: Septicemia (except in labor) | 405,898 | 21.42 | 21.87 | 2.1 |
| 3: Bacterial infection; unspecified site | 1,625 | 18.22 | 18.71 | 2.7 |
| 4: Mycoses | 10,755 | 27.56 | 28.10 | 2.0 |
| 5: HIV infection | 7,495 | 26.99 | 27.66 | 2.5 |
| 6: Hepatitis | 10,121 | 35.23 | 36.29 | 3.0 |
| 7: Viral infection | 14,033 | 16.01 | 16.46 | 2.8 |
| 8: Other infections; including parasitic | 4,117 | 13.46 | 14.09 | 4.7 |
| 9: Sexually transmitted infections (not HIV or hepatitis) | 699 | 14.16 | 14.45 | 2.0 |
| 10: Immunizations and screening for infectious disease | 149 | 14.09 | 14.09 | 0.0 |
| 11: Cancer of head and neck | 8,915 | 20.43 | 21.40 | 4.8 |
| 12: Cancer of esophagus | 3,956 | 26.69 | 27.68 | 3.7 |
| 13: Cancer of stomach | 7,171 | 23.79 | 24.68 | 3.8 |
| 14: Cancer of colon | 37,987 | 17.16 | 17.53 | 2.2 |
| 15: Cancer of rectum and anus | 12,496 | 21.26 | 21.68 | 2.0 |
| 16: Cancer of liver and | 5,935 | 24.08 | 25.14 | 4.4 |

| | | | 30-Day Readmission Rate per 100 Index Events | | |
|---|------------------------------|---|---|--|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States | |
| intrahepatic bile duct | | | | | |
| 17: Cancer of pancreas | 12,641 | 24.41 | 25.74 | 5.5 | |
| 18: Cancer of other GI organs; peritoneum | 7,043 | 24.00 | 25.27 | 5.3 | |
| 19: Cancer of bronchus; lung | 51,885 | 20.46 | 21.12 | 3.2 | |
| 20: Cancer; other respiratory and intrathoracic | 952 | 17.65 | 19.85 | 12.5 | |
| 21: Cancer of bone and connective tissue | 3,007 | 20.02 | 20.82 | 4.0 | |
| 22: Melanomas of skin | 989 | 13.25 | 13.65 | 3.0 | |
| 23: Other non-epithelial cancer of skin | 2,483 | 15.99 | 16.31 | 2.0 | |
| 24: Cancer of breast | 18,788 | 9.27 | 9.41 | 1.5 | |
| 25: Cancer of uterus | 10,304 | 11.81 | 12.35 | 4.5 | |
| 26: Cancer of cervix | 1,616 | 20.92 | 21.78 | 4.1 | |
| 27: Cancer of ovary | 6,450 | 21.83 | 22.71 | 4.0 | |
| 28: Cancer of other female genital organs | 2,155 | 15.82 | 16.15 | 2.0 | |
| 29: Cancer of prostate | 24,323 | 6.93 | 7.13 | 3.0 | |
| 31: Cancer of other male genital organs | 308 | 22.73 | 23.38 | 2.9 | |
| 32: Cancer of bladder | 14,780 | 22.18 | 23.03 | 3.8 | |
| 33: Cancer of kidney and renal pelvis | 14,717 | 13.23 | 13.84 | 4.6 | |
| 34: Cancer of other urinary organs | 1,716 | 16.14 | 16.96 | 5.1 | |
| 35: Cancer of brain and nervous system | 7,670 | 20.38 | 21.51 | 5.6 | |
| 36: Cancer of thyroid | 3,775 | 9.35 | 9.48 | 1.4 | |
| 37: Hodgkin`s disease | 584 | 32.36 | 33.39 | 3.2 | |

| | | 30-Day Read per 100 Inc | mission Rate lex Events | Percentage Increase in |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 38: Non-Hodgkin`s Iymphoma | 12,102 | 34.35 | 35.44 | 3.2 |
| 39: Leukemias | 8,871 | 33.62 | 34.86 | 3.7 |
| 40: Multiple myeloma | 5,748 | 28.97 | 29.91 | 3.2 |
| 41: Cancer; other and unspecified primary | 1,514 | 16.84 | 17.90 | 6.3 |
| 42: Secondary malignancies | 65,482 | 23.08 | 23.97 | 3.8 |
| 43: Malignant neoplasm without specification of site | 2,667 | 24.03 | 24.82 | 3.3 |
| 44: Neoplasms of unspecified nature or uncertain behavior | 14,836 | 25.74 | 26.54 | 3.1 |
| 45: Maintenance chemotherapy; radiotherapy | 25,281 | 61.29 | 61.98 | 1.1 |
| 46: Benign neoplasm of uterus | 4,909 | 7.29 | 7.37 | 1.1 |
| 47: Other and unspecified benign neoplasm | 34,259 | 12.74 | 13.18 | 3.4 |
| 48: Thyroid disorders | 9,288 | 10.81 | 11.00 | 1.8 |
| 49: Diabetes mellitus without complication | 3,189 | 16.09 | 16.31 | 1.4 |
| 50: Diabetes mellitus with complications | 159,421 | 23.81 | 24.31 | 2.1 |
| 51: Other endocrine disorders | 23,041 | 20.09 | 20.52 | 2.1 |
| 52: Nutritional deficiencies | 4,402 | 24.49 | 24.97 | 1.9 |
| 53: Disorders of lipid metabolism | 147 | 14.29 | 14.29 | 0.0 |
| 54: Gout and other crystal arthropathies | 9,186 | 17.46 | 17.92 | 2.6 |
| 55: Fluid and electrolyte disorders | 192,425 | 20.33 | 20.81 | 2.4 |

| | | 30-Day Read per 100 Inc | mission Rate lex Events | Percentage Increase in |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 56: Cystic fibrosis | 1,507 | 20.31 | 20.77 | 2.3 |
| 57: Immunity disorders | 306 | 28.76 | 30.07 | 4.5 |
| 58: Other nutritional; endocrine; and metabolic disorders | 33,388 | 16.81 | 17.25 | 2.6 |
| 59: Deficiency and other anemia | 92,656 | 23.19 | 23.74 | 2.4 |
| 60: Acute posthemorrhagic anemia | 16,095 | 21.09 | 21.66 | 2.7 |
| 61: Sickle cell anemia | 15,360 | 34.26 | 35.22 | 2.8 |
| 62: Coagulation and hemorrhagic disorders | 9,731 | 27.73 | 28.53 | 2.9 |
| 63: Diseases of white blood cells | 17,437 | 26.37 | 27.06 | 2.6 |
| 64: Other hematologic conditions | 1,256 | 22.05 | 22.93 | 4.0 |
| 76: Meningitis (except that caused by tuberculosis or sexually transmitted disease) | 3,339 | 16.17 | 17.07 | 5.6 |
| 77: Encephalitis (except that caused by tuberculosis or sexually transmitted disease) | 1,945 | 17.22 | 18.46 | 7.2 |
| 78: Other CNS infection and poliomyelitis | 2,052 | 22.86 | 23.64 | 3.4 |
| 79: Parkinson`s disease | 6,647 | 13.25 | 13.44 | 1.4 |
| 80: Multiple sclerosis | 6,730 | 13.80 | 14.28 | 3.4 |
| 81: Other hereditary and degenerative nervous system conditions | 17,788 | 16.29 | 16.83 | 3.3 |
| 82: Paralysis | 1,732 | 17.21 | 17.55 | 2.0 |
| 83: Epilepsy; convulsions | 70,843 | 15.62 | 16.10 | 3.1 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|--|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 84: Headache; including migraine | 12,433 | 13.04 | 13.56 | 4.0 |
| 85: Coma; stupor; and brain damage | 7,730 | 16.96 | 17.50 | 3.2 |
| 87: Retinal detachments; defects; vascular occlusion; and retinopathy | 1,091 | 9.90 | 10.36 | 4.6 |
| 88: Glaucoma | 274 | 13.14 | 13.50 | 2.8 |
| 89: Blindness and vision defects | 1,943 | 12.30 | 13.02 | 5.9 |
| 90: Inflammation; infection of eye (except that caused by tuberculosis or sexually transmitted disease) | 3,104 | 13.24 | 13.85 | 4.6 |
| 91: Other eye disorders | 1,835 | 11.55 | 12.15 | 5.2 |
| 92: Otitis media and related conditions | 1,410 | 11.49 | 12.13 | 5.6 |
| 93: Conditions associated with dizziness or vertigo | 29,165 | 8.27 | 8.53 | 3.2 |
| 94: Other ear and sense organ disorders | 1,672 | 13.22 | 13.64 | 3.2 |
| 95: Other nervous system disorders | 88,536 | 18.37 | 18.94 | 3.1 |
| 96: Heart valve disorders | 47,909 | 21.73 | 23.16 | 6.6 |
| 97: Peri-; endo-; and myocarditis; cardiomyopathy (except that caused by tuberculosis or sexually transmitted disease) | 19,317 | 21.89 | 22.71 | 3.8 |
| 98: Essential hypertension | 21,797 | 11.97 | 12.30 | 2.8 |
| 99: Hypertension with complications and secondary hypertension | 101,943 | 24.08 | 24.58 | 2.1 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 100: Acute myocardial infarction | 205,949 | 19.66 | 20.41 | 3.8 |
| 101: Coronary atherosclerosis and other heart disease | 220,481 | 15.44 | 15.99 | 3.5 |
| 102: Nonspecific chest pain | 148,512 | 13.50 | 14.00 | 3.7 |
| 103: Pulmonary heart disease | 67,061 | 16.71 | 17.20 | 2.9 |
| 104: Other and ill-defined heart disease | 3,266 | 13.38 | 14.15 | 5.7 |
| 105: Conduction disorders | 29,215 | 11.42 | 11.86 | 3.9 |
| 106: Cardiac dysrhythmias | 345,225 | 16.73 | 17.27 | 3.3 |
| 107: Cardiac arrest and ventricular fibrillation | 3,236 | 18.88 | 19.96 | 5.7 |
| 108: Congestive heart failure; nonhypertensive | 468,709 | 25.05 | 25.70 | 2.6 |
| 109: Acute cerebrovascular disease | 232,510 | 14.73 | 15.23 | 3.4 |
| 110: Occlusion or stenosis of precerebral arteries | 59,225 | 10.67 | 10.97 | 2.8 |
| 111: Other and ill-defined cerebrovascular disease | 7,927 | 11.30 | 11.86 | 4.9 |
| 112: Transient cerebral ischemia | 80,396 | 11.31 | 11.66 | 3.1 |
| 113: Late effects of cerebrovascular disease | 8,073 | 15.88 | 16.31 | 2.7 |
| 114: Peripheral and visceral atherosclerosis | 77,210 | 19.15 | 19.65 | 2.6 |
| 115: Aortic; peripheral; and visceral artery aneurysms | 35,223 | 15.16 | 15.95 | 5.2 |
| 116: Aortic and peripheral arterial embolism or | 11,656 | 22.50 | 23.07 | 2.6 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| thrombosis | | | | |
| 117: Other circulatory disease | 61,972 | 17.52 | 17.98 | 2.6 |
| 118: Phlebitis; thrombophlebitis and thromboembolism | 67,493 | 16.60 | 17.07 | 2.8 |
| 119: Varicose veins of lower extremity | 1,061 | 16.21 | 16.68 | 2.9 |
| 120: Hemorrhoids | 12,573 | 17.41 | 17.74 | 1.9 |
| 121: Other diseases of veins and lymphatics | 9,833 | 20.34 | 21.06 | 3.5 |
| 122: Pneumonia (except that caused by tuberculosis or sexually transmitted disease) | 443,388 | 18.25 | 18.69 | 2.4 |
| 123: Influenza | 11,775 | 11.77 | 12.03 | 2.2 |
| 124: Acute and chronic tonsillitis | 809 | 7.17 | 7.54 | 5.2 |
| 125: Acute bronchitis | 26,446 | 13.00 | 13.32 | 2.4 |
| 126: Other upper respiratory infections | 9,847 | 13.01 | 13.48 | 3.6 |
| 127: Chronic obstructive pulmonary disease and bronchiectasis | 356,562 | 21.99 | 22.40 | 1.9 |
| 128: Asthma | 85,354 | 18.61 | 18.99 | 2.0 |
| 129: Aspiration pneumonitis; food/vomitus | 95,063 | 20.93 | 21.22 | 1.4 |
| 130: Pleurisy; pneumothorax; pulmonary collapse | 37,794 | 25.02 | 25.81 | 3.1 |
| 131: Respiratory failure; insufficiency; arrest (adult) | 136,502 | 24.58 | 25.19 | 2.5 |
| 132: Lung disease due to external agents | 2,365 | 20.25 | 21.02 | 3.8 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 133: Other lower respiratory disease | 44,901 | 18.94 | 19.54 | 3.1 |
| 134: Other upper respiratory disease | 12,106 | 18.43 | 19.09 | 3.6 |
| 135: Intestinal infection | 86,677 | 22.59 | 23.04 | 2.0 |
| 136: Disorders of teeth and jaw | 3,436 | 12.34 | 12.75 | 3.3 |
| 137: Diseases of mouth; excluding dental | 5,261 | 14.94 | 15.32 | 2.5 |
| 138: Esophageal disorders | 41,912 | 16.29 | 16.70 | 2.5 |
| 139: Gastroduodenal ulcer (except hemorrhage) | 12,983 | 17.04 | 17.62 | 3.4 |
| 140: Gastritis and duodenitis | 34,213 | 19.14 | 19.57 | 2.2 |
| 141: Other disorders of stomach and duodenum | 26,484 | 24.64 | 25.32 | 2.8 |
| 142: Appendicitis and other appendiceal conditions | 18,638 | 9.55 | 9.84 | 3.0 |
| 143: Abdominal hernia | 62,904 | 13.38 | 13.74 | 2.6 |
| 144: Regional enteritis and ulcerative colitis | 16,244 | 23.15 | 24.02 | 3.7 |
| 145: Intestinal obstruction without hernia | 131,147 | 17.46 | 17.93 | 2.7 |
| 146: Diverticulosis and diverticulitis | 107,020 | 14.55 | 14.91 | 2.5 |
| 147: Anal and rectal conditions | 13,837 | 16.30 | 16.68 | 2.3 |
| 148: Peritonitis and intestinal abscess | 8,092 | 27.39 | 28.28 | 3.2 |
| 149: Biliary tract disease | 107,986 | 14.87 | 15.38 | 3.4 |
| 151: Other liver diseases | 36,907 | 32.88 | 34.08 | 3.7 |
| 152: Pancreatic disorders | 66,802 | 19.80 | 20.58 | 4.0 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| (not diabetes) | | | | |
| 153: Gastrointestinal hemorrhage | 155,410 | 18.75 | 19.27 | 2.8 |
| 154: Noninfectious gastroenteritis | 41,796 | 16.12 | 16.57 | 2.8 |
| 155: Other gastrointestinal disorders | 73,673 | 20.39 | 20.92 | 2.6 |
| 156: Nephritis; nephrosis; renal sclerosis | 1,488 | 25.00 | 25.54 | 2.2 |
| 157: Acute and unspecified renal failure | 228,134 | 22.04 | 22.60 | 2.5 |
| 158: Chronic kidney disease | 8,075 | 26.13 | 26.81 | 2.6 |
| 159: Urinary tract infections | 277,674 | 18.31 | 18.66 | 1.9 |
| 160: Calculus of urinary tract | 32,158 | 13.21 | 13.53 | 2.4 |
| 161: Other diseases of kidney and ureters | 12,302 | 19.05 | 19.64 | 3.1 |
| 162: Other diseases of bladder and urethra | 8,649 | 18.28 | 18.82 | 3.0 |
| 163: Genitourinary symptoms and ill-defined conditions | 15,706 | 20.16 | 20.69 | 2.6 |
| 164: Hyperplasia of prostate | 23,248 | 11.51 | 11.75 | 2.1 |
| 165: Inflammatory conditions of male genital organs | 6,873 | 13.90 | 14.22 | 2.3 |
| 166: Other male genital disorders | 2,842 | 15.94 | 16.50 | 3.5 |
| 167: Nonmalignant breast conditions | 2,997 | 13.38 | 13.65 | 2.0 |
| 168: Inflammatory diseases of female pelvic organs | 3,024 | 15.18 | 15.68 | 3.3 |
| 169: Endometriosis | 940 | 7.98 | 8.19 | 2.7 |
| 170: Prolapse of female | 25,446 | 3.53 | 3.66 | 3.7 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|--|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| genital organs | | | | |
| 171: Menstrual disorders | 3,085 | 9.17 | 9.37 | 2.1 |
| 172: Ovarian cyst | 2,420 | 10.29 | 10.50 | 2.0 |
| 173: Menopausal disorders | 2,132 | 12.48 | 12.81 | 2.6 |
| 175: Other female genital disorders | 7,410 | 11.86 | 12.27 | 3.4 |
| 177: Spontaneous abortion | 119 | 6.72 | 6.72 | 0.0 |
| 180: Ectopic pregnancy | 159 | 8.18 | 8.18 | 0.0 |
| 181: Other complications of pregnancy | 3,919 | 20.31 | 20.80 | 2.4 |
| 182: Hemorrhage during pregnancy; abruptio placenta; placenta previa | 266 | 17.67 | 18.80 | 6.4 |
| 183: Hypertension complicating pregnancy; childbirth and the puerperium | 1,544 | 16.78 | 16.90 | 0.8 |
| 184: Early or threatened labor | 1,068 | 23.69 | 24.06 | 1.6 |
| 185: Prolonged pregnancy | 665 | 2.11 | 2.26 | 7.2 |
| 186: Diabetes or abnormal glucose tolerance complicating pregnancy; childbirth; or the puerperium | 822 | 19.34 | 19.83 | 2.5 |
| 187: Malposition; malpresentation | 414 | 4.59 | 4.59 | 0.0 |
| 188: Fetopelvic disproportion; obstruction | 153 | 4.58 | 5.23 | 14.3 |
| 189: Previous C-section | 2,229 | 3.99 | 4.08 | 2.3 |
| 190: Fetal distress and abnormal forces of labor | 584 | 4.62 | 4.62 | 0.0 |
| 191: Polyhydramnios and | 821 | 8.53 | 8.77 | 2.9 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|--|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| other problems of amniotic cavity | | | | |
| 192: Umbilical cord complication | 421 | 1.43 | 1.43 | 0.0 |
| 193: OB-related trauma to perineum and vulva | 1,159 | 1.12 | 1.12 | 0.0 |
| 195: Other complications of birth; puerperium affecting management of mother | 2,716 | 8.51 | 8.62 | 1.3 |
| 196: Normal pregnancy and/or delivery | 450 | 1.56 | 1.56 | 0.0 |
| 197: Skin and subcutaneous tissue infections | 178,097 | 16.09 | 16.45 | 2.2 |
| 198: Other inflammatory condition of skin | 2,901 | 21.06 | 21.68 | 2.9 |
| 199: Chronic ulcer of skin | 25,685 | 21.13 | 21.62 | 2.3 |
| 200: Other skin disorders | 2,589 | 17.69 | 18.00 | 1.7 |
| 201: Infective arthritis and osteomyelitis (except that caused by tuberculosis or sexually transmitted disease) | 24,731 | 19.24 | 19.77 | 2.8 |
| 202: Rheumatoid arthritis and related disease | 5,487 | 12.96 | 13.30 | 2.7 |
| 203: Osteoarthritis | 363,295 | 5.57 | 5.70 | 2.2 |
| 204: Other non-traumatic joint disorders | 16,187 | 11.90 | 12.27 | 3.1 |
| 205: Spondylosis; intervertebral disc disorders; other back problems | 178,614 | 10.62 | 10.93 | 2.9 |
| 206: Osteoporosis | 146 | 21.23 | 21.92 | 3.2 |
| 207: Pathological fracture | 31,718 | 19.03 | 19.46 | 2.3 |
| 208: Acquired foot | 1,705 | 7.57 | 7.68 | 1.5 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| deformities | | | | |
| 209: Other acquired deformities | 17,245 | 9.92 | 10.22 | 3.0 |
| 210: Systemic lupus erythematosus and connective tissue disorders | 3,920 | 26.79 | 27.37 | 2.2 |
| 211: Other connective tissue disease | 47,933 | 13.59 | 14.00 | 3.0 |
| 212: Other bone disease and musculoskeletal deformities | 21,751 | 12.33 | 12.63 | 2.5 |
| 213: Cardiac and circulatory congenital anomalies | 2,502 | 13.75 | 14.71 | 7.0 |
| 214: Digestive congenital anomalies | 713 | 15.71 | 16.69 | 6.3 |
| 215: Genitourinary congenital anomalies | 1,201 | 16.24 | 16.57 | 2.1 |
| 216: Nervous system congenital anomalies | 397 | 17.88 | 18.39 | 2.8 |
| 217: Other congenital anomalies | 4,654 | 8.10 | 8.36 | 3.2 |
| 225: Joint disorders and dislocations; trauma-related | 6,439 | 11.14 | 11.34 | 1.8 |
| 226: Fracture of neck of femur (hip) | 172,566 | 13.89 | 14.19 | 2.2 |
| 227: Spinal cord injury | 2,420 | 17.69 | 19.09 | 7.9 |
| 228: Skull and face fractures | 7,091 | 11.90 | 12.38 | 4.0 |
| 229: Fracture of upper limb | 43,161 | 11.92 | 12.33 | 3.5 |
| 230: Fracture of lower limb | 61,816 | 13.59 | 13.96 | 2.7 |
| 231: Other fractures | 87,204 | 14.05 | 14.50 | 3.2 |
| 232: Sprains and strains | 9,057 | 11.52 | 11.91 | 3.4 |
| 233: Intracranial injury | 55,231 | 16.40 | 17.16 | 4.6 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 234: Crushing injury or internal injury | 13,358 | 15.29 | 15.98 | 4.5 |
| 235: Open wounds of head; neck; and trunk | 5,616 | 12.29 | 12.84 | 4.5 |
| 236: Open wounds of extremities | 5,507 | 13.78 | 14.11 | 2.4 |
| 237: Complication of device; implant or graft | 274,669 | 21.23 | 21.83 | 2.8 |
| 238: Complications of surgical procedures or medical care | 155,584 | 20.50 | 21.08 | 2.9 |
| 239: Superficial injury; contusion | 16,861 | 15.88 | 16.23 | 2.2 |
| 240: Burns | 4,110 | 16.40 | 17.54 | 7.0 |
| 241: Poisoning by psychotropic agents | 14,978 | 12.82 | 13.27 | 3.5 |
| 242: Poisoning by other medications and drugs | 27,934 | 15.48 | 15.99 | 3.3 |
| 243: Poisoning by nonmedicinal substances | 2,659 | 11.51 | 12.19 | 5.9 |
| 244: Other injuries and conditions due to external causes | 19,846 | 15.79 | 16.34 | 3.5 |
| 245: Syncope | 95,001 | 11.90 | 12.29 | 3.2 |
| 246: Fever of unknown origin | 17,447 | 19.54 | 20.14 | 3.1 |
| 247: Lymphadenitis | 1,297 | 18.50 | 19.20 | 3.8 |
| 248: Gangrene | 17,575 | 33.79 | 34.34 | 1.6 |
| 249: Shock | 554 | 23.11 | 24.19 | 4.7 |
| 250: Nausea and vomiting | 16,042 | 22.97 | 23.48 | 2.2 |
| 251: Abdominal pain | 36,919 | 20.14 | 20.89 | 3.7 |
| 252: Malaise and fatigue | 14,952 | 16.77 | 17.19 | 2.5 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 253: Allergic reactions | 5,477 | 15.45 | 15.94 | 3.2 |
| 254: Rehabilitation care; fitting of prostheses; and adjustment of devices | 1,558 | 11.04 | 11.10 | 0.6 |
| 256: Medical examination/evaluation | 101 | 22.77 | 23.76 | 4.3 |
| 257: Other aftercare | 1,889 | 14.08 | 14.45 | 2.6 |
| 258: Other screening for suspected conditions (not mental disorders or infectious disease) | 522 | 17.43 | 17.63 | 1.1 |
| 259: Residual codes; unclassified | 43,683 | 18.57 | 19.14 | 3.0 |
| 650: Adjustment disorders | 1,361 | 13.45 | 13.89 | 3.3 |
| 651: Anxiety disorders | 4,677 | 15.72 | 16.02 | 1.9 |
| 652: Attention-deficit | 245 | 19.59 | 20.00 | 2.1 |
| 653: Delirium | 37,650 | 14.28 | 14.72 | 3.1 |
| 654: Developmental disorders | 613 | 19.09 | 19.41 | 1.7 |
| 655: Disorders usually diagnosed in infancy | 115 | 15.65 | 17.39 | 11.1 |
| 656: Impulse control disorders | 417 | 17.03 | 17.51 | 2.8 |
| 657: Mood disorders | 45,648 | 18.81 | 19.38 | 3.0 |
| 658: Personality disorders | 431 | 25.52 | 26.45 | 3.6 |
| 659: Schizophrenia and other psychotic disorders | 47,114 | 22.04 | 22.37 | 1.5 |
| 660: Alcohol-related disorders | 30,228 | 20.72 | 21.24 | 2.5 |
| 661: Substance-related disorders | 32,424 | 18.68 | 19.28 | 3.2 |

| | | 30-Day Read per 100 Inc | Percentage Increase in | |
|---|------------------------------|---|---|--|
| Clinical Classifications Software Principal Diagnosis Category for the Index Event | Number of Index Events | Readmissions Limited to the Same State as the Index Event | Readmissions Considered from Any State | the Readmission Rates When Patients Were Followed Across States |
| 662: Suicide and intentional self-inflicted injury | 240 | 16.67 | 17.50 | 5.0 |
| 663: Screening and history of mental health and substance abuse codes | 11,302 | 28.35 | 29.03 | 2.4 |
| 670: Miscellaneous disorders | 3,592 | 13.34 | 14.23 | 6.7 |